

Product Datasheet

WT1 Antibody (WT1/857 + 6F-H2) NBP2-44607-0.1mg

Unit Size: 0.1 mg

Store at 4C.

www.novusbio.com



technical@novusbio.com

Publications: 8

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-44607

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-44607



NBP2-44607-0.1mg

WT1 Antibody (WT1/857 + 6F-H2)

Product Information	
Unit Size	0.1 mg
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	WT1/857 + 6F-H2
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa/IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA


Product Description	
Description	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-47858) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	7490
Gene Symbol	WT1
Species	Human, Mouse, Rat
Reactivity Notes	Use in Mouse reported in scientific literature (PMID:33631230).
Marker	Wilm's Tumor & Mesothelial Marker
Specificity/Sensitivity	Recognizes a 47-55kDa-tumor suppressor protein, identified as Wilms tumor and mesothelioma. WT1 protein has been identified in proliferative mesothelial cells, malignant mesothelioma, ovarian carcinoma, gonadoblastoma, nephroblastoma, and desmoplastic small round cell tumor. Lung adenocarcinomas rarely stain positive with this antibody. WT1 protein expression in mesothelial cells has become a reliable marker for the diagnosis of mesotheliomas.
Immunogen	Recombinant full length human WT1 protein (WT1/857); Recombinant fragment aa1-181 of human WT1 (6F-H2) (Uniprot: P19544)

Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin 1-2 ug/ml
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.




Images


Immunohistochemistry-Paraffin: WT1 Antibody (WT1/857 + 6F-H2) [NBP2-44607] - Formalin-fixed, paraffin-embedded human Mesothelioma stained with WT1 Antibody (WT1/857 + 6F-H2).



Immunohistochemistry-Paraffin: WT1 Antibody (WT1/857 + 6F-H2) [NBP2-44607] - Formalin-fixed, paraffin-embedded Rat kidney stained with WT1 monoclonal antibody (WT1/857+6F-H2).



Immunohistochemistry-Paraffin: WT1 Antibody (WT1/857 + 6F-H2) [NBP2-44607] - Formalin-fixed, paraffin-embedded human Fetal kidney stained with WT1 Monoclonal Antibody (WT1/857+6F-H2).



Publications

Jillian Williquett, Chantal Allamargot, Hua Sun AMPK-SP1–Guided Dynein Expression Represents a New Energy-Responsive Mechanism and Therapeutic Target for Diabetic Nephropathy *Kidney360* 2024-04-01 [PMID: 38467599]

Wang Z, Wu Q, Wang H et al. Diosgenin protects against podocyte injury in early phase of diabetic nephropathy through regulating SIRT6 *Phytomedicine : international journal of phytotherapy and phytopharmacology* 2022-06-13 [PMID: 35728388] (IF/IHC, Mouse)

Yang X, Luo W, Li L et al. CDK9 inhibition improves diabetic nephropathy by reducing inflammation in the kidneys *Toxicology and applied pharmacology* 2021-02-22 [PMID: 33631230] (IHC-P, Mouse)

Zheng C, Huang L et al. Inhibition of STAT3 in tubular epithelial cells prevents kidney fibrosis and nephropathy in STZ-induced diabetic mice. *Cell Death Dis* 2019-07-11 [PMID: 31699972] (IF/IHC, Mouse)

Liang G, Song L et al. Fibroblast growth factor 1 ameliorates diabetic nephropathy by an anti-inflammatory mechanism. *Kidney Int* 2018-01-01 [PMID: 28750927] (IF/IHC, Human)

Luo W, Chen X, Ye L et al. Kaempferol attenuates streptozotocin-induced diabetic nephropathy by downregulating TRAF6 expression: the role of TRAF6 in diabetic nephropathy *Journal of Ethnopharmacology* 2020-11-01 [PMID: 33152432] (IHC-P, Mouse)

Chen X, Zhang X, Xu J et Al. AZD4547 Attenuates Lipopolysaccharide-Induced Acute Kidney Injury by Inhibiting Inflammation: The Role of FGFR1 in Renal Tubular Epithelial Cells *Drug Des Devel Ther* 2020-02-26 [PMID: 32161443] (IHC-P, IF/IHC, Mouse)

Chen X, Zhao Y, Xu J et al. The Nephroprotective Effect of TNF Receptor-Associated Factor 6 (TRAF6) Blockade on LPS-Induced Acute Renal Injury Through the Inhibition if Inflammation and Oxidative Stress *Med. Sci. Monit.* 2020-01-13 [PMID: 31929494] (IHC-P, Mouse)



**Novus Biologicals USA**

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-44607-0.1mg

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
H00007490-Q01-10ug	Recombinant Human WT1 GST (N-Term) Protein
292-G2-050	IGF-II/IGF2 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-44607

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

