Product Datasheet

Chromogranin A Antibody (LK2H10 + PHE5) - IHC-Prediluted NBP2-44771

Unit Size: 7 ml

Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-44771

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-44771



NBP2-44771


Chromogranin A Antibody (LK2H10 + PHE5) - IHC-Prediluted

Product Information Init Size 7 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C. Clonality Monoclonal Clone LK2H10 + PHE5 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, ph 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific appl	Chromogranin A Antibody (LK2H10 + PHE5) - IHC-Prediluted	
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C. Clonality Monoclonal Clone LK2H10 + PHE5 Preservative 0.05% Sodium Azide Isotype IlgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Information	
Storage Store at 4C. Clonality Monoclonal Clone LK2H10 + PHE5 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Product Description Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Application Notes Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml	Unit Size	7 ml
Clone	Concentration	·
Clone LK2H10 + PHE5 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Product Description Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 10.25 - 0.5 ug/ml Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Storage	Store at 4C.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Product Description Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
Isotype	Clone	LK2H10 + PHE5
Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Product Description Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	0.05% Sodium Azide
Buffer 10 mM PBS with 0.05% BSA Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG1 Kappa
Product Description The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. Host Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS with 0.05% BSA
titration; the antibody is ready-to-use and optimized for staining. Mouse Gene ID 1113 Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	
Gene Symbol CHGA Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Mouse
Species Human, Mouse, Rat, Porcine, Monkey Marker Neuroendocrine Marker Immunogen Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	1113
MarkerNeuroendocrine MarkerImmunogenHuman pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645)Product Application DetailsImmunohistochemistry, Immunohistochemistry-ParaffinApplicationsImmunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	CHGA
Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645) Product Application Details	Species	Human, Mouse, Rat, Porcine, Monkey
Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	Neuroendocrine Marker
Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	Human pheochromocytoma (LK2H10 & PHE5) (Uniprot: P10645)
Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Application Notes Immunohistochemistry (Formalin-fixed): 0.1-0.2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml
	Application Notes	of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.




Images

Immunohistochemistry-Paraffin: Chromogranin A Antibody (LK2H10 + PHE5) - IHC-Prediluted [NBP2-44771] - Formalin-fixed, paraffinembedded human small cell lung carcinoma stained with Chromogranin A MAb (LK2H10+PHE5)



Immunohistochemistry-Paraffin: Chromogranin A Antibody (LK2H10 + PHE5) - IHC-Prediluted [NBP2-44771] - Formalin-fixed paraffinembedded human Adrenal Gland stained with Chromogranin A Monoclonal Antibody (PHE5)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-44771

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP2-61453-10ug Recombinant Human Chromogranin A His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-44771

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

