Product Datasheet

dedicator of cytokinesis 8 Antibody (OTI4H10) NBP2-46471

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-46471

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-46471



NBP2-46471


dedicator of cytokinesis 8 Antibody (OTI4H10)

dedicator of cytokinesis 8 Antibody (OTI4H10)	
Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	OTI4H10
Preservative	0.02% Sodium Azide
Isotype	lgG1
Purity	Immunogen affinity purified
Buffer	PBS (pH 7.3), 1.0% BSA and 50% Glycerol
Target Molecular Weight	238.3 kDa
Product Description	
Description	Novus Biologicals Mouse dedicator of cytokinesis 8 Antibody (OTI4H10) (NBP2-46471) is a monoclonal antibody validated for use in IHC and WB. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	81704
Gene Symbol	DOCK8
Species	Human, Mouse, Rat
Reactivity Notes	Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.
Immunogen	Human recombinant protein fragment corresponding to amino acids 832-1160 of human DOCK8 (NP_982272) produced in E.coli.
Product Application Details	

	` , .
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry
Recommended Dilutions	Western Blot 1:2000, Immunohistochemistry 1:150, Immunohistochemistry-




Western Blot: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY dedicator of cytokinesis 8.







Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human tonsil tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Adenocarcinoma of Human breast tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human colon tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)



Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Adenocarcinoma of Human colon tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human Kidney tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Carcinoma of Human kidney tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human liver tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Carcinoma of Human liver tissue.(Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human lung tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Carcinoma of Human lung tissue .(Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human Ovary tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Adenocarcinoma of Human ovary tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human pancreas tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Carcinoma of Human pancreas tissue. (Heatinduced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human thyroid tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Carcinoma of Human thyroid tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human endometrium tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Adenocarcinoma of Human endometrium tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human prostate tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Carcinoma of Human prostate tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human bladder tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)




Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Carcinoma of Human bladder tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)



Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human lymph node tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)



Immunohistochemistry: dedicator of cytokinesis 8 Antibody (4H10) [NBP2-46471] - Analysis of Human lymphoma tissue. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120C for 3min)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-46471

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)

NBP3-17049PEP dedicator of cytokinesis 8 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-46471

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

