# **Product Datasheet**

# TFF1/pS2 Antibody (SPM573) NBP2-32900-0.1mg

Unit Size: 0.1 mg Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-32900

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-32900



## NBP2-32900-0.1mg

TFF1/pS2 Antibody (SPM573)

| TFF1/pS2 Antibody (SPM573)         |  |
|------------------------------------|--|
| Product Information                |  |
| Unit Size                          | 0.1 mg   |
| Concentration                      | 0.2 mg/ml  |
| Storage                            | Store at 4C.   |
| Clonality                          | Monoclonal   |
| Clone                              | SPM573   |
| Preservative                       | 0.05% Sodium Azide   |
| Isotype                            | IgG1 Kappa   |
| Purity                             | Protein A or G purified  |
| Buffer                             | 10 mM PBS with 0.05% BSA   |
| Target Molecular Weight            | 6.5 kDa  |
| <b>Product Description</b>         |  |
| Description                        | 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34806)  Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.   |
| Host                               | Mouse  |
| Gene ID                            | 7031   |
| Gene Symbol                        | TFF1   |
| Species                            | Human, Cynomolgus Monkey   |
| Specificity/Sensitivity            | It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy. |
| Immunogen                          | Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human TFF1/pS2. (Uniprot: P04155)   |
| <b>Product Application Details</b> |  |
| Applications                       | Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin   |
| Recommended Dilutions              | Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml  |
| 1                                  |  |




#### **Application Notes**

Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

#### **Images**

Immunohistochemistry-Paraffin: TFF1/pS2 Antibody (SPM573) [NBP2-32900] - Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with TFF1/pS2 Antibody (SPM573).





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP2-32900-0.1mg

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

NBP2-35042-5ug Recombinant Human TFF1/pS2 Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-32900

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

