# **Product Datasheet**

## CD45RB Antibody (BRA-11 (same as BRA-11G)) NBP2-33021-0.1mg

Unit Size: 0.1 mg

Store at 4C.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-33021

Updated 7/16/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-33021



## NBP2-33021-0.1mg

CD45RB Antibody (BRA-11 (same as BRA-11G))

Product Information     Unit Size   0.1 mg     Concentration   0.2 mg/ml     Storage   Store at 4C.     Clonality   Monoclonal     Clone   BRA-11 (same as BRA-11G)     Preservative   0.05% Sodium Azide     Isotype   IgG1 Kappa     Purity   Protein A or G purfied     Buffer   10 mM PBS with 0.05% BSA     Product Description   200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G.     Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available   WITHOUT BSA & azide at 1.0 mg/ml (NBP2-34564)     Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to = s0c.   s0c.     Host   Mouse   Gene ID     Gene Bymbol   PTPRC   Species     Species   Human, Monkey (Negative), Rat (Negative)   Reactivity Notes     Boc.Oll Marker   B-Cell Marker     Immunogen   Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)     Product Application Details   Mestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Pararafin     Recommended Di		
Concentration 0.2 mg/ml   Storage Stor at 4C.   Clonality Monoclonal   Clone BRA-11 (same as BRA-11G)   Preservative 0.05% Sodium Azide   Isotype IgG1 Kappa   Purity Protein A or G purified   Buffer 10 mM PBS with 0.05% BSA   Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)   Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.   Host Mouse   Gene ID 5788   Gene Symbol PTPRC   Species Human, Monkey (Negative), Rat (Negative)   Reactivity Notes Does not react with Rat or Rhesus Monkey.   Marker B-Cell Marker   Immunohistochemistry. Immunofuorescence, Immunohistochemistry. Immunofluorescence, Immunohistochemistry. Immunofluorescence 1-2 ug/ml, Immunohistochemi	Product Information	
Storage Store at 4C.   Clonality Monoclonal   Clone BRA-11 (same as BRA-11G)   Preservative 0.05% Sodium Azide   Isotype IgG1 Kappa   Purity Protein A or G purified   Buffer 10 mM PBS with 0.05% BSA   Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)   Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.   Host Mouse   Gene ID 5788   Gene Symbol PTPRC   Species Human, Monkey (Negative), Rat (Negative)   Reactivity Notes Does not react with Rat or Rhesus Monkey.   Marker B-Cell Marker   Immunogen Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)   Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunoliuorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Inmunocytochemistry/ Immunofluorescence 1-2 ug/ml, Im	Unit Size	0.1 mg
Clonality   Monoclonal     Clone   BRA-11 (same as BRA-11G)     Preservative   0.05% Sodium Azide     Isotype   IgG1 Kappa     Purity   Protein A or G purified     Buffer   10 mM PBS with 0.05% BSA     Product Description   200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)     Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.     Host   Mouse     Gene ID   5788     Gene Symbol   PTPRC     Species   Human, Monkey (Negative), Rat (Negative)     Reactivity Notes   Does not react with Rat or Rhesus Monkey.     Marker   B-Cell Marker     Immunogen   Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)     Product Application Details   Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry/ Immunohistochemistry/ Immunohistochemistry/ Immunohistochemistry, Immunohiston tot.	Concentration	0.2 mg/ml
Clone BRA-11 (same as BRA-11G)   Preservative 0.05% Sodium Azide   Isotype IgG1 Kappa   Purity Protein A or G purified   Buffer 10 mM PBS with 0.05% BSA   Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)   Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.   Host Mouse   Gene ID 5788   Gene Symbol PTPRC   Species Human, Monkey (Negative), Rat (Negative)   Reactivity Notes Does not react with Rat or Rhesus Monkey.   Marker B-Cell Marker   Immunogen No-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)   Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry-Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin   Recommended Dilutions Western Blot, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry for 20 minutes at RT. Staining of tormalin-fixed tissues requires heating tissue sections in 10m Tris with 1mM Tis with 1mM	Storage	Store at 4C.
Preservative   0.05% Sodium Azide     Isotype   IgG1 Kappa     Purity   Protein A or G purified     Buffer   10 mM PBS with 0.05% BSA     Product Description   200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WiTHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)     Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.     Host   Mouse     Gene ID   5788     Gene Symbol   PTPRC     Species   Human, Monkey (Negative), Rat (Negative)     Reactivity Notes   Does not react with Rat or Rhesus Monkey.     Marker   B-Cell Marker     Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, 2 ug/ml, Immunohistochemistry-Paraffin     Recommended Dilutions   Western Blot, Flow Cytometry, Involocytochemistry/Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed; 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed itsues requires heating tissue sections in 10m Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
Isotype IgG1 Kappa   Purity Protein A or G purified   Buffer 10 mM PBS with 0.05% BSA   Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)   Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.   Host Mouse   Gene ID 5788   Gene Symbol PTPRC   Species Human, Monkey (Negative), Rat (Negative)   Reactivity Notes Does not react with Rat or Rhesus Monkey.   Marker B-Cell Marker   Immunohistochemistry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Flow Cytometry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemist	Clone	BRA-11 (same as BRA-11G)
Purity   Protein A or G purified     Buffer   10 mM PBS with 0.05% BSA     Product Description   200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)     Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.     Host   Mouse     Gene ID   5788     Gene Symbol   PTPRC     Species   Human, Monkey (Negative), Rat (Negative)     Reactivity Notes   Does not react with Rat or Rhesus Monkey.     Marker   B-Cell Marker     Immunogen   Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)     Product Application Details   Western Blot, Flow Cytometry, Immunocytochemistry-Paraffin     Recommended Dilutions   Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry-Paraffin     Replication Notes   Immunohistochemistry, Formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	0.05% Sodium Azide
Buffer 10 mM PBS with 0.05% BSA   Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)   Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.   Host Mouse   Gene ID 5788   Gene Symbol PTPRC   Species Human, Monkey (Negative), Rat (Negative)   Reactivity Notes Does not react with Rat or Rhesus Monkey.   Marker B-Cell Marker   Immunogen Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)   Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry-Paraffin   Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry-Paraffin   Recommended Dilutions Western Blot, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry -Paraffin   Application Notes Immunohistochemistry, Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10m Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG1 Kappa
Product Description   200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)     Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.     Host   Mouse     Gene ID   5788     Gene Symbol   PTPRC     Species   Human, Monkey (Negative), Rat (Negative)     Reactivity Notes   Does not react with Rat or Rhesus Monkey.     Marker   B-Cell Marker     Immunogen   Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)     Product Application Details   Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin     Recommended Dilutions   Western Blot, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Oromalin-fixed): 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
Description200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID5788Gene SymbolPTPRCSpeciesHuman, Monkey (Negative), Rat (Negative)Reactivity NotesDoes not react with Rat or Rhesus Monkey.MarkerB-Cell MarkerImmunogenNon-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Inmunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry 1-2 ug/ml flor 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS with 0.05% BSA
Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostMouseGene ID5788Gene SymbolPTPRCSpeciesHuman, Monkey (Negative), Rat (Negative)Reactivity NotesDoes not react with Rat or Rhesus Monkey.MarkerB-Cell MarkerImmunogenNon-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry (Formalin 1-2 ug/mlApplication NotesWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed issues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
-80C.   Host Mouse   Gene ID 5788   Gene Symbol PTPRC   Species Human, Monkey (Negative), Rat (Negative)   Reactivity Notes Does not react with Rat or Rhesus Monkey.   Marker B-Cell Marker   Immunogen Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)   Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin   Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry-Paraffin 1-2 ug/ml   Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34564)
Gene ID5788Gene SymbolPTPRCSpeciesHuman, Monkey (Negative), Rat (Negative)Reactivity NotesDoes not react with Rat or Rhesus Monkey.MarkerB-Cell MarkerImmunogenNon-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)Product Application DetailsApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.		
Gene Symbol PTPRC   Species Human, Monkey (Negative), Rat (Negative)   Reactivity Notes Does not react with Rat or Rhesus Monkey.   Marker B-Cell Marker   Immunogen Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)   Product Application Details Applications   Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin   Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml   Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Mouse
SpeciesHuman, Monkey (Negative), Rat (Negative)Reactivity NotesDoes not react with Rat or Rhesus Monkey.MarkerB-Cell MarkerImmunogenNon-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	5788
Reactivity Notes Does not react with Rat or Rhesus Monkey.   Marker B-Cell Marker   Immunogen Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)   Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin   Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml   Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	PTPRC
Marker B-Cell Marker   Immunogen Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)   Product Application Details Applications   Mestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin   Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml   Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human, Monkey (Negative), Rat (Negative)
ImmunogenNon-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)Product Application DetailsApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Reactivity Notes	Does not react with Rat or Rhesus Monkey.
Product Application DetailsApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	B-Cell Marker
ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)
Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	
of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml,
	Application Notes	of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.

www.novusbio.com




#### Images

Western Blot: CD45RB Antibody (BRA-11 (same as BRA-11G)) [NBP2-33021] - Western Blot Analysis of human Raji cell lysate using CD45RB Antibody (BRA-11 (same as BRA-11G)).

Flow Cytometry: CD45RB Antibody (BRA-11 (same as BRA-11G))

Mouse- IgG-CF488 (Blue); Isotype Control (Red).

tonsil stained with CD45RB Ab (BRA-11).



www.novusbio.com





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-33021-0.1mg

210-TA-005	TNF-alpha [Unconjugated]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-33021

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

