

Product Datasheet

Ku70/XRCC6 Antibody (KU729)

NBP2-34247-0.1mg

Unit Size: 0.1 mg

Store at 4C.

www.novusbio.com



technical@novusbio.com

Reviews: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-34247

Updated 7/16/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-34247



NBP2-34247-0.1mg


Ku70/XRCC6 Antibody (KU729)

Product Information	
Unit Size	0.1 mg
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	KU729
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Product Description	
Description	<p>200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-34663)</p> <p>Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.</p>
Host	Mouse
Gene ID	2547
Gene Symbol	XRCC6
Species	Human, Primate, Chicken (Negative), Mouse (Negative), Rat (Negative)
Reactivity Notes	Does not react with Mouse, Rat and Chicken.
Marker	Nuclear Marker
Specificity/Sensitivity	Recognizes a dimer of two proteins of 70kDa and ~80kDa, identified as two subunits of Ku. This monoclonal antibody recognizes a conformational epitope of p70/p80 dimer, which is destroyed during Western blotting. The p70/p80 dimer is important for function of a 460kDa DNA-dependent protein kinase. Ku protein plays a role in cell signaling, proliferation, DNA repair, replication, transcriptional activation, and apoptosis.
Immunogen	Nuclear extract of human HL-60 cells
Product Application Details	
Applications	Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Flow Cytometry 1-2 ug/million cells, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Flow (Intracellular)
Application Notes	Immunocytochemistry (Acetone-fixed cells): 0.5-1.0ug/ml for 30 minutes at RT. Optimal dilution for a specific application should be determined. Optimal dilution for a specific application should be determined.




Images


Immunocytochemistry/Immunofluorescence: Ku70/XRCC6 Antibody (KU729) [NBP2-34247] - Immunofluorescent analysis of HeLa cells using Ku70/XRCC6 antibody (KU729). Image from verified customer review.




Flow Cytometry: Ku70/XRCC6 Antibody (KU729) [NBP2-34247] - Flow Cytometric Analysis of human Ku (p70/p80) on 293T cells. Black: cells alone; Grey: Isotype Control; Green: AF488-labeled Ku70/XRCC6 Antibody (KU729).




Immunocytochemistry/Immunofluorescence: Ku70/XRCC6 Antibody (KU729) [NBP2-34247] - Immunofluorescent staining of paraformaldehyde-fixed HeLa cells using followed by goat anti-Mouse IgG-CF488 (Green). Phalloidin (Red) is used to label cell membrane.




Flow Cytometry: Ku70/XRCC6 Antibody (KU729) [NBP2-34247] - Flow Cytometry of human Ku (p70/p80) on K562 cells. Black: cells alone; Green: Isotype Control; Red: PE-labeled Ku70/XRCC6 Antibody (KU729).



Flow (Intracellular): Ku70/XRCC6 Antibody (KU729) [NBP2-34247] - An intracellular stain was performed on Hek293 cells with Ku70/XRCC6 Antibody (KU729) NBP2-34663AF647 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.



Flow (Intracellular): Ku70/XRCC6 Antibody (KU729) [NBP2-34247] - An intracellular stain was performed on Hek293 cells with Ku70/XRCC6 Antibody (KU729) NBP2-34663AF647 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-34247-0.1mg

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
H00002547-P02-2ug	Recombinant Human Ku70/XRCC6 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34247

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

