

# Product Datasheet

## Cytokeratin 19 Antibody (BA17) NBP2-34271-0.1mg

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-34271](http://www.novusbio.com/NBP2-34271)

Updated 1/14/2021 v.20.1

**Earn rewards for product reviews and publications.**

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-34271](http://www.novusbio.com/reviews/destination/NBP2-34271)



**NBP2-34271-0.1mg**

Cytokeratin 19 Antibody (BA17)

**Product Information**

<b>Unit Size</b>	0.1 mg
<b>Concentration</b>	1.0 mg/ml
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Monoclonal
<b>Clone</b>	BA17
<b>Preservative</b>	0.02% Sodium Azide
<b>Isotype</b>	IgG1 Kappa
<b>Purity</b>	Protein A purified
<b>Buffer</b>	PBS
<b>Target Molecular Weight</b>	40 kDa

**Product Description**

<b>Host</b>	Mouse
<b>Gene ID</b>	3880
<b>Gene Symbol</b>	KRT19
<b>Species</b>	Human
<b>Marker</b>	Pancreatic Stem Cell Marker
<b>Specificity/Sensitivity</b>	This Ab reacts with the rod domain of human cytokeratin 19 (CK19), a polypeptide of 40kDa. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.
<b>Immunogen</b>	Human mammary epithelial organoids

**Product Application Details**


<b>Applications</b>	Western Blot, Simple Western, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Immunoprecipitation, SDS-Page
<b>Recommended Dilutions</b>	Western Blot 0.5-1ug/ml, Simple Western 10 ug/ml, Flow Cytometry 0.5-1ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2ug/ml, Immunoprecipitation 1-2ug/500ug protein, Immunohistochemistry-Paraffin 0.5-1ug/ml, Immunohistochemistry-Frozen 0.5-1ug/ml, SDS-Page
<b>Application Notes</b>	ELISA (For coating, order NBP2-34560 (BSA free version of NBP2-34271)). The 7mL size is a pre-diluted size and no additional dilutions are required before using this item for the intended application.

## Images

Immunohistochemistry-Paraffin: Cytokeratin 19 Antibody (BA17) [NBP2-34271] - Formalin-fixed, paraffin-embedded human colon stained with Cytokeratin 19 MAb (BA17)




Flow Cytometry: Cytokeratin 19 Antibody (BA17) [NBP2-34271] - An intracellular stain was performed on HepG2 cells with Cytokeratin 19 antibody (BA17) NBP2-34560APC (blue) and a matched isotype control NBP2-27287APC (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Allophycocyanin.




Simple Western: Cytokeratin 19 Antibody (BA17) [NBP2-34271] - Simple Western lane view shows a specific band for Cytokeratin 19 in 0.2 mg/ml of HeLa lysate(s). This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Simple Western: Cytokeratin 19 Antibody (BA17) [NBP2-34271] - Electropherogram image of the corresponding Simple Western lane. Cytokeratin 19 antibody was used at 10 ug/ml dilution of HeLa lysates(s) respectively.



## SDS-Page: Cytokeratin 19 Antibody (BA17) [NBP2-34271]





## Novus Biologicals USA

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

## Bio-Techne Canada

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

## Bio-Techne Ltd

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

## General Contact Information

[www.novusbio.com](http://www.novusbio.com)  
Technical Support: [nb-technical@bio-techne.com](mailto:nb-technical@bio-techne.com)  
Orders: [nb-customerservice@bio-techne.com](mailto:nb-customerservice@bio-techne.com)  
General: [novus@novusbio.com](mailto:novus@novusbio.com)

## Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-34271](http://www.novusbio.com/reviews/submit/NBP2-34271)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)