Product Datasheet

ZAP70 Antibody (2F3.2) [DyLight 405] NBP2-47774V

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47774V

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-47774V



NBP2-47774V

ZAP70 Antibody (2F3.2) [DyLight 405]

| Product Information | 2/11 /0 /1111100dy (21 3.2) [DyEig | 11. 400] |
|---|------------------------------------|---|
| Please see the vial label for concentration. If unlisted please contact technical services. | Product Information | |
| Storage Store at 4C in the dark. Clonality Monoclonal Clone 2F3.2 Preservative 0.05% Sodium Azide Isotype IgG2a Kappa Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7535 Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with jumutated CLL, samples, whereas no band is observed in the Ig-mutated CLL samples. In Western blotting of cell lysates of CD19-positive Purified leukemia cells from patients with E1-tymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of AG31 cells (carcinoma cell line), no band is observed in the Ig-mutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-tymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of AG31 cells (carcinoma cell line), no band is observed in the Ig-mutated CLL samples. In Western blotting of cell lysates of AG31 cells (carcinoma cell line), no band is observed in the Ig-mutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombiant ZAP70 positive Including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Unit Size | 0.1 ml |
| Clone 2F3.2 Preservative 0.05% Sodium Azide Isotype 1gG2a Kappa Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7535 Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated CLL, the antibody labels a band corresponding to XeP70. In Western blotting of cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated CLL samples. In Western blotting of cell lysates of A31 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 protein is expressed in leukemia (CLL) cases as well. Anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Concentration | · |
| Clone 2F3.2 Preservative 0.05% Sodium Azide Isotype IgG2a Kappa Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7535 Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated CLL samples. In Western blotting of cell visates of CL12 samples. In Western blotting of cell visates of A431 cells (carcinoma cell line), no band is observed. ZAP70 protein in expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) casses as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (CLL) cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Storage | Store at 4C in the dark. |
| Preservative D.0.5% Sodium Azide | Clonality | Monoclonal |
| IgG2a Kappa | Clone | 2F3.2 |
| Conjugate DyLight 405 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7535 Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of control of this protein translation is via the IgVH gene. In Western blotting of of CD19-positive Purified leukemia cells from patients with Ig-unmutated and Ig-mutated CLL, the antibody labels a band corresponding to ZAP70. In western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band corresponding to ZaP70 protein in expression is an | Preservative | 0.05% Sodium Azide |
| Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Host Mouse Gene ID 7535 Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell lysates of cD19-positive Purified leukemia cells from patients with Ig-muntated and Ig-mutated CLL, the antibody labels a band corresponding to ZAP70 in the Ig-mutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples, whereas no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 propative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Isotype | IgG2a Kappa |
| Buffer SomM Sodium Borate | Conjugate | DyLight 405 |
| Product Description This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 7535 Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70 in the Igunumutated CLL, the antibody labels a band corresponding to ZAP70 in the Igunumutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band corresponding to SAP70 in the Igunumutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band corresponding to SAP70 in the Igunumutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of Ad31 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Igunmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Purity | Protein A or G purified |
| This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 7535 Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated and Igmutated CLL, the antibody labels a band corresponding to ZAP70 in the Ig-unmutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of A431 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Buffer | 50mM Sodium Borate |
| Volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Mouse Gene ID 7535 Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated and Ig-mutated CLL, the antibody labels a band corresponding to ZAP70 in the Ig-unmutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of A431 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Product Description | |
| Gene Symbol ZAP70 Species Human Marker Chronic Lymphocytic Leukemia Marker Specificity/Sensitivity ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated and Igmutated CLL, the antibody labels a band corresponding to ZAP70 in the Ig-unmutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of A431 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Description | volume of this product. The volume will be greater than or equal to the unit size |
| Species Human | Host | Mouse |
| Species Human | Gene ID | 7535 |
| Marker Chronic Lymphocytic Leukemia Marker ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell Iysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell Iysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated and Igmutated CLL, the antibody labels a band corresponding to ZAP70 in the Ig-unmutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples. In Western blotting of cell Iysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell Iysates of A431 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Gene Symbol | ZAP70 |
| ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. In Western blotting of whole cell Iysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell Iysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated and Igmutated CLL, the antibody labels a band corresponding to ZAP70 in the Ig-unmutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of A431 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Species | Human |
| Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated and Igmutated CLL, the antibody labels a band corresponding to ZAP70 in the Igunmutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of A431 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis. Immunogen Recombinant ZAP-70 protein including residues 1-254 and encompassing SH2 domains of human ZAP70 (Uniprot: P43403) | Marker | Chronic Lymphocytic Leukemia Marker |
| domains of human ZAP70 (Uniprot: P43403) Notes | Specificity/Sensitivity | Control of this protein translation is via the IgVH gene. In Western blotting of whole cell lysates of normal peripheral blood mononuclear cells, the antibody labels a band corresponding to ZAP70. In Western blotting of whole cell lysates of CD19-positive Purified leukemia cells from patients with Ig-unmutated and Igmutated CLL, the antibody labels a band corresponding to ZAP70 in the Ig-unmutated CLL samples, whereas no band is observed in the Ig-mutated CLL samples. In Western blotting of cell lysates of Jurkat cells (T-lymphoblastic cell line), the antibody labels a band of 70kDa protein. In Western blotting of cell lysates of A431 cells (carcinoma cell line), no band is observed. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases |
| | Immunogen | |
| Product Application Details | | DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. |

Product Application Details

Simple Western, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, **Applications** Immunofluorescence



| | Simple Western, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready |
|-------------------|---|
| Application Notes | Optimal dilution of this antibody should be experimentally determined. |





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-47774V

NBP1-96981V Mouse IgG2a Kappa Isotype Control (M2AK) [DyLight 405]

NBP1-87000PEP ZAP70 Recombinant Protein Antigen

202-IL-010 IL-2 [Unconjugated] 3709-KS-010 ZAP70 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47774V

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

