## **Product Datasheet**

### Androgen R/NR3C4 Antibody (SPM335) [DyLight 550] NBP2-47855R

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47855R

Updated 10/26/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-47855R



#### NBP2-47855R

Androgen R/NR3C4 Antibody (SPM335) [DyLight 550]

services.   Storage Store at 4C in the dark.   Clonality Monocional   Clone SPM335   Preservative 0.05% Sodium Azide   lsotype IgG1 Kappa   Conjugate DyLight 550   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor. It doe cross react with storgen, progesterone, or gluccoortical receptors. The expression of AR is reportedly inversely cortelated with histlogic grade i.e. J.   Specificity/Sensitivity Recognizes a avortein of 110kDa, which is identified as androgen receptor. It doe cross react with all energy anti-androgen therapy. Anti-androgen these proposed, as a marke hormone-responsiveness and thus it may be useful indentifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful cinically in difff	5	
Concentration   Please see the vial label for concentration. If unlisted please contact technical services.     Storage   Store at 4C in the dark.     Clonality   Monoclonal     Clone   SPM335     Preservative   0.05% Sodium Azide     Isotype   IgG1 Kappa     Conjugate   DyLight 550     Purity   Protein A or G purified     Buffer   50mM Sodium Borate     Product Description   This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.     Host   Mouse     Gene ID   367     Gene Symbol   AR     Species   Human, Mouse (Negative)     Reactivity Notes   Does not react with Mouse.     Marker   Marker of Androgen Dependence     Specificity/Sensitivity   Recognizes a protein of 110kDa, which is identified as androgen receptor. (A differentiated prostate currors. Ner Abas been proposed, as a mark bornore. responsiveness and thus it may be useful indentifying patients like benefit from anti-androgen therapy. Anti-androgen receptor, it doe cross react with estrogen, progesterone, or gluccocricid receptors. The expression of AR is reported/parafitin tissues.     Immunogen   A syntheti	Product Information	
services.   Storage Store at 4C in the dark.   Clonality Monoclonal   Clone SPM335   Preservative 0.05% Sodium Azide   lsotype IgG1 Kappa   Conjugate DyLight 550   Purity Protein A or G purified   Buffer Som Mosodium Borate   Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker of Androgen Dependence Specificity/Sensitivity   Recognizes a protein of 110kDa, which is identified as androgen receptor. It doe cross react with estrogen, progesterone, or glucocorticol receptors. The explicitity/Sensitivity   Recognizes a solution of thus reported runors show higher expression of AR is reported runors. In prostate cancer. AR has been proposed, as a marke homone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor tha been useful cinically in differentiated prostate cancer. An has been proposed, as a marke homone-responsiveness and thus it may be useful in identifying patients like bene	Unit Size	0.1 ml
Clonality Monoclonal   Clone SPM335   Preservative 0.05% Sodium Azide   Isotype IgG1 Kappa   Conjugate DyLight 550   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor. The expression of AR is reportedly inversely correlated with histologic grade i.e. desmoplastic timos. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness show higher expression than the poorly. differentiated tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor. The expression of restate tumors. This sonoclonal antibody i superb for staining of formalin/paraffit tissues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Clone SPM335   Preservative 0.05% Sodium Azide   Isotype IgG1 Kappa   Conjugate DyLight 550   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description Description   Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor. It does care react with full length, and the newly described A form of the receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e., differentiated turnors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in dentifying patients like semplastic inchoepithelinoma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.   Immunogen A synthetic peptide, as 302-318 (STEDTAEYSPFKGGYTK) of human Androgen Preceptor stabe on the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide   Isotype IgG1 Kappa   Conjugate DyLight 550   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor (A differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marke benefit from anti-androgen therapy. Anti-androgen receptor has been useful ulcinically in differentiated tumors. In prostate cancer, AR has been useful ulcinically in differentiated tumors. Show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marke benefit from anti-androgen therapy. Anti-androgen receptor has been useful ulcinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplasic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.	Clonality	Monoclonal
Isotype IgG1 Kappa   Conjugate DyLight 550   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor. It does cross react with estrogen, progesterone, or glucocriticoi receptors. The very soin of AR is reportedly inversely correlated with histologic grade i.e. v differentiated trumors. In prostate cancer, AR has been useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor more as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor may be a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen freezof as a ab2-318 (STEDTAEYSPFKGGYTK) of human Androgen Lawred (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher	Clone	SPM335
Conjugate DyLight 550   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor. It does cross react with full length, and the newly described A form of the receptor. It does cross react with storgen, progesterone, or gluccorticold receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. ufifferentiated trons the prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morphaeform basal cell carcinoma (mRCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody is superb for staining of formalin/paraffin tissues.   Immunogen A synthetic peptide, as 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarid Product Application	Preservative	0.05% Sodium Azide
Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor. It doe cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen tereptor. It does cross react trith esting of formalin/parafit itssues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Androg   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie   Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence exercementers and intreacting andifferencementers anditreacting andifferenc	Isotype	IgG1 Kappa
Buffer 50mM Sodium Borate   Product Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor (A reacts with full length, and the newly described A form of the receptor. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful indentifying patients like benefit from anti-androgen threapy. Anti-androgen receptor Ans bene useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody is superb for statining of formalin/parafit tissues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiatient immunofilorescence, Immunohistochemistry, Immunofluorescence, Immunohistochemistry.Paraffin, CyTOF-ready, Immunofluorescence	Conjugate	DyLight 550
Product Description   This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.     Host   Mouse     Gene ID   367     Gene Symbol   AR     Species   Human, Mouse (Negative)     Reactivity Notes   Does not react with Mouse.     Marker   Marker of Androgen Dependence     Specificity/Sensitivity   Recognizes a protein of 110kDa, which is identified as androgen receptor (A reacts with full length, and the newly described A form of the receptors. The expression of AR is reported) inversely correlated with histologic grade i.e. v differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors show higher expression than the poorly differentiated prostate tumors and whis it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichcoeptitelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.     Immunogen   A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)     Notes   DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie     Product Application Details   Flow Cytometry, Immunocytochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Purity	Protein A or G purified
Description This conjugate is made on demand. Actual recovery may vary from the state volume of this product. The volume will be greater than or equal to the unit si stated on the datasheet.   Host Mouse   Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor (A reacts with full length, and the newly described A form of the receptor. It does cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful circically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie   Product Application Details Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry.Paraffin, CyTOF-ready, Immunofluoresc	Buffer	50mM Sodium Borate
Volume of this product. The volume will be greater than or equal to the unit sistated on the datasheet.HostMouseGene ID367Gene SymbolARSpeciesHuman, Mouse (Negative)Reactivity NotesDoes not react with Mouse.MarkerMarker of Androgen DependenceSpecificity/SensitivityRecognizes a protein of 110kDa, which is identified as androgen receptor. It doe cross react with full length, and the newly described A form of the receptor. It doe cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated prostate tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.ImmunogenA synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarieProduct Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence	Product Description	
Gene ID 367   Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor (A reacts with full length, and the newly described A form of the receptor. It does cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaried for Cytometry, Immunocytochemistry-Paraffin, CyTOF-ready, Immunofluorescence, Immunohistochemistry. Immunofluorescence, Immunohistochemistry. Immunofluorescence	Description	This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Gene Symbol AR   Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor. It does cross react with full length, and the newly described A form of the receptor. It does cross react with full length, and the newly described A form of the receptor. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated trumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiated trumors. In prostate cancor, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaried   Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Host	Mouse
Species Human, Mouse (Negative)   Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor (A reacts with full length, and the newly described A form of the receptor. It does cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaried   Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Gene ID	367
Reactivity Notes Does not react with Mouse.   Marker Marker of Androgen Dependence   Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor (A reacts with full length, and the newly described A form of the receptor. It does cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie   Product Application Details Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry. Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Gene Symbol	AR
Marker   Marker of Androgen Dependence     Specificity/Sensitivity   Recognizes a protein of 110kDa, which is identified as androgen receptor (A reacts with full length, and the newly described A form of the receptor. It does cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. with full length and the newly described A form of the receptor. It does cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. with full rentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.     Immunogen   A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)     Notes   DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie     Product Application Details   Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Species	Human, Mouse (Negative)
Specificity/Sensitivity Recognizes a protein of 110kDa, which is identified as androgen receptor (A reacts with full length, and the newly described A form of the receptor. It does cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marke hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.   Immunogen A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)   Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie   Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence	Reactivity Notes	Does not react with Mouse.
reacts with full length, and the newly described A form of the receptor. It does cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. v differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marked hormone-responsiveness and thus it may be useful in identifying patients like benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody i superb for staining of formalin/paraffin tissues.ImmunogenA synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Andro R/NR3C4 (Uniprot: P10275)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie Immunohistochemistry, Immunohytochemistry/Immunofluorescence, Immunohistochemistry, Immunohytochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Marker	Marker of Androgen Dependence
R/NR3C4 (Uniprot: P10275)   Notes   DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries   Product Application Details   Applications   Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Specificity/Sensitivity	expression of AR is reportedly inversely correlated with histologic grade i.e. well differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marker of hormone-responsiveness and thus it may be useful in identifying patients likely to benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This monoclonal antibody is
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie   Product Application Details   Applications   Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Immunogen	A synthetic peptide, aa 302-318 (STEDTAEYSPFKGGYTK) of human Androgen R/NR3C4 (Uniprot: P10275)
Product Application Details   Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Notes	
Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence		DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence	Product Application Details	
		Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
	Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready



Application Notes	Optimal dilution of this antibody should be experimentally determined.
-------------------	--

www.novusbio.com





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-47855R

NBP1-43319R	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 550]
H00000367-Q01-10ug	Recombinant Human Androgen R/NR3C4 GST (N-Term) Protein
236-EG-200	EGF [Unconjugated]
NBP2-69854	Human Androgen R/NR3C4 ELISA Kit (Chemiluminescence)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47855R

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

