## **Product Datasheet**

# Hepatocyte Specific Antigen Antibody (HepPar1) - IHC-Prediluted NBP2-48068

Unit Size: 7 ml

Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-48068

Updated 6/14/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-48068




#### NBP2-48068

| Hepatocyte Specific Antigen Antibody (HepPar1) - IHC-Prediluted  |  |
|--|--|
| Product Information  |  |
| 7 ml   |  |
| Please see the vial label for concentration. If unlisted please contact technical services.  |  |
| Store at 4C.   |  |
| Monoclonal   |  |
| HepPar1  |  |
| 0.05% Sodium Azide   |  |
| IgG1 Kappa   |  |
| Protein A or G purified  |  |
| 10 mM PBS with 0.05% BSA   |  |
| Product Description  |  |
| The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining.  |  |
| Mouse  |  |
| Human, Mouse, Canine   |  |
| Mouse reactivity reported in scientific literature (PMID: 27542264).   |  |
| Hepatocellular Marker  |  |
| Extract of a formalin-fixed, rejected-allograft of a human liver   |  |
| Product Application Details  |  |
| Immunohistochemistry, Immunohistochemistry-Paraffin  |  |
| Immunohistochemistry, Immunohistochemistry-Paraffin  |  |
| Immunohistochemistry-Paraffin 0.25 - 0.5 ug/ml for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10 - 20 min followed by cooling at RT for 20 minutes. The 7mL size is a pre-diluted size and no additional dilutions are required before using this item for the intended application. |  |
|  |  |



## **Images**

Immunohistochemistry-Paraffin: Hepatocyte Specific Antigen Antibody (HepPar1) - IHC-Prediluted [NBP2-48068] - Formalin-fixed, paraffinembedded human Liver stained with Hepatocyte Specific Antigen Antibody (HepPar1).



Immunohistochemistry-Paraffin: Hepatocyte Specific Antigen Antibody (HepPar1) - IHC-Prediluted [NBP2-48068] - Human Hepatocellular Carcinoma stained with HepPar-1 Monoclonal Antibody (HepPar1).





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP2-48068**

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-48068

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

