# **Product Datasheet**

# Neuro2a Chloroquine Treated / Untreated Cell Lysate NBP2-49688

Unit Size: 2 Vials

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

**Publications: 1** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-49688

Updated 11/28/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-49688



# NBP2-49688


Neuro2a Chloroquine Treated / Untreated Cell Lysate

| Nedroza Officioquine Treated / Officialed Octi Lysate |   |
|---|---|
| Product Information                                   |   |
| Unit Size   | 2 Vials   |
| Concentration   | 1.0 mg/ml   |
| Storage   | Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.  |
| Preservative  | No Preservative   |
| Buffer  | 1X Laemmli Sample Buffer [31.5 mM Tris HCl, 1% SDS, 10% Glycerol, 0.005% bromophenol blue, pH 6.8]  |
| Product Description                                   |   |
| Description   | NBP2-49688 contains 1 vial each of 0.1ml of Neuro2a Chloroquine treated and 0.1ml of Neuro2a untreated cell lysate. This lysate set is useful as a positive and negative controls for targets such as LC3, p62 etc. The researchers should use anti-mouse primary antibodies when working with these lysates.   |
| Species   | Mouse   |
| Marker  | Autophagy/LC3 Positive Control  |
| Preparation Method                                    | Neuro2A cells were cultured under standard laboratory condition until semi-confluent (70-80%). The cells were then treated with or without Chloroquine to 50 uM for 24 hours. The cells were washed in PBS and directly lysed into 1x Laemmli sample buffer containing BME. Each lysate is sonicated and boiled before being tested in Western blot for reactivity to LC3. Tubulin reactivity in each lysate is shown as a loading control. |
| Kit Components  | 0.1 ml Neuro2a Chloroquine treated whole cell lysate, 0.1 ml Neuro2a untreated whole cell lysate  |
| Lysate Type   | Cell  |
| Lysate Tissue   | Brain   |
| Lysate Subcellular Fraction                           | Chloroquine Treated / Untreated   |
| Product Application Details                           |   |
| Applications  | Western Blot, SDS-Page  |
| Recommended Dilutions                                 | Western Blot, SDS-Page  |
| Application Notes                                     | Neuro2a Chloroquine treated / untreated lysates are provided as positive and negative control for Western blot analysis in Autophagy research. The lysates are provided as one vial of treated and one vial of untreated samples. Use 10 ul per lane for a standard mini-gel blot (approx. 1 mg/ml).  |
|   |   |




#### **Images**

Western Blot: Neuro2a Chloroquine Treated / Untreated Cell Lysate [NBP2-49688] - Mouse Neuoblast cells (Neuro2A) were treated with (+) and without (-) 50 uM Chloroquine overnight. Whole cell protein lysates were prepared in 1x Laemmli sample buffer and approximately 10 ug of each lysate (NBP2-49688) was separated on a 4-15% gel by SDS-PAGE, transferred to 0.2 um PVDF membrane and blocked in 5% nonfat milk in TBST. The membrane was probed with 1 ug/ml anti-LC3 (NB100-2220) and 1 ug/ml anti-alpha tubulin (NB100-690) as a loading control, and detected with the appropriate secondary antibodies using chemiluminescence.



Western Blot: Neuro2a Chloroquine Treated / Untreated Cell Lysate [NBP2-49688] - Western Blot: [NBP2-49688] - 10 ug of Chloroquine treated (+) and untreated (-) Neuro2A lysates in 1x Laemmli sample buffer (NBP2-49688) were separated on a 4-15% gel by SDS-PAGE, transferred to 0.2 um PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 1 ug/ml monoclonal anti-LC3 (NBP2--46892) and detected with an anti-rabbit secondary antibody using chemiluminescence.



#### **Publications**

Cachon-Gonzalez Mb, Wang S, Cox Tm Expression of Ripk1 and DAM genes correlates with severity and progression of Krabbe disease Human molecular genetics 2021-06-25 [PMID: 34172992] (WB)



# Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

# Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-49688

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

