## **Product Datasheet**

# CD43/Sialophorin Antibody (SPN/1766R) - Azide and BSA Free NBP2-54344-100ug

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54344

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-54344




## NBP2-54344-100ug

| CD43/Sialophorin Antibody (SPN/1766R) - Azide and BSA Free |   |
|--|---|
| Product Information  |   |
| Unit Size  | 100 ug  |
| Concentration  | 1 mg/ml   |
| Storage  | Store at -20 to -80C. Avoid freeze-thaw cycles.   |
| Clonality  | Monoclonal  |
| Clone  | SPN/1766R   |
| Preservative   | No Preservative   |
| Isotype  | IgG   |
| Purity   | Protein A or G purified   |
| Buffer   | 10 mM PBS   |
| Product Description  |   |
| Description  | 1.0 mg/ml of recombinant monoclonal antibody purified by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53288).  Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.  |
| Host   | Rabbit  |
| Gene ID  | 6693  |
| Gene Symbol  | SPN   |
| Species  | Human   |
| Marker   | T-Cell Marker   |
| Specificity/Sensitivity                                    | It recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma. |
| Immunogen  | Recombinant full-length human CD43/Sialophorin protein (Uniprot: P16150)  |
| Product Application Details                                |   |
| Applications   | Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready   |
| Recommended Dilutions                                      | Western Blot, Flow Cytometry 0.5 - 1 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1 - 2 ug/ml, Immunohistochemistry-Paraffin 0.5 - 1.0 ug/ml, CyTOF-ready  |
| Application Notes  | Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.   |




#### **Images**


Western Blot: CD43/Sialophorin Antibody (SPN/1766R) - Azide and BSA Free [NBP2-54344] - Western Blot Analysis of K562 cell lysate using CD43/Sialophorin Antibody (SPN/1766R).




Immunocytochemistry/Immunofluorescence: CD43/Sialophorin Antibody (SPN/1766R) - Azide and BSA Free [NBP2-54344] - Immunofluorescence Analysis of K562 cells labeling CD43 with CD43/Sialophorin Antibody (SPN/1766R) followed by Goat anti-Rabbit-IgG-CF488 (Green). The nuclear counterstain is NucSpot .



Immunohistochemistry-Paraffin: CD43/Sialophorin Antibody (SPN/1766R) - Azide and BSA Free [NBP2-54344] - Formalin-fixed, paraffin-embedded human Tonsil stained with CD43 Recombinant Rabbit Monoclonal Antibody (SPN/1766R).



Flow Cytometry: CD43/Sialophorin Antibody (SPN/1766R) - Azide and BSA Free [NBP2-54344] - Flow Cytometric Analysis of PFA-fixed K562 cells using CD43/Sialophorin Antibody (SPN/1766R) followed by Goat anti-Rabbit- IgG-CF488 (Blue); Isotype Control (Red)





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## Products Related to NBP2-54344-100ug

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-23472 Recombinant Human CD43/Sialophorin His Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54344

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

