Product Datasheet

CD3 epsilon Antibody (PC3/188A) [Alexa Fluor® 488] NBP2-54405AF488

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54405AF488

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-54405AF488



NBP2-54405AF488

CD3 epsilon Antibody (PC3/188A) [Alexa Fluor® 488]

Product Information	CD3 epsilon Antibody (PC3/166A) [Alexa Fidol® 466]	
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone PC3/188A Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic.	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone PC3/188A Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage, In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Unit Size	100 ul
Clone PC3/188A Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely asociated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Concentration	·
Clone PC3/188A Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Storage	Store at 4C in the dark.
Preservative Isotype IgG1 Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Clonality	Monoclonal
IgG1	Clone	PC3/188A
Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Isotype	IgG1
Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Conjugate	Alexa Fluor 488
Product Description Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Purity	Protein A or G purified
Host Mouse Gene ID 916 Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Buffer	50mM Sodium Borate
Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Product Description	
Gene Symbol CD3E Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Reactivity Notes Others not known. Marker T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Host	Mouse
Species Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit Others not known. T-Cell Marker Specificity/Sensitivity Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Gene ID	916
Reactivity Notes Others not known. T-Cell Marker Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Gene Symbol	CD3E
T-Cell Marker Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Species	Human, Rat, Porcine, Bovine, Chicken, Equine, Guinea Pig, Rabbit
Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Reactivity Notes	Others not known.
chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms. Immunogen A synthetic peptide corresponding to amino acids 156-168 of the cytoplasmic	Marker	T-Cell Marker
	Specificity/Sensitivity	chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.
	Immunogen	



Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, CyTOF-ready, Immunohistochemistry (Negative)
Recommended Dilutions	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry (Negative), CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Images

CD3 epsilon Antibody (PC3/188A) [Alexa Fluor® 488]







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-54405AF488

DDXCM01A488 Mouse IgG1 Isotype Control [Alexa Fluor® 488]
NBP2-22752 Recombinant Human CD3 epsilon His Protein

210-TA-005 TNF-alpha [Unconjugated] 9850-CD-050 CD3 epsilon [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54405AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

