## **Product Datasheet**

### PAX8 Antibody (PAX8/1491) [Alexa Fluor® 647] NBP2-54486AF647

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54486AF647

Updated 10/26/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54486AF647



#### NBP2-54486AF647

PAX8 Antibody (PAX8/1491) [Alexa Fluor® 647]

Product InformationUnit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePAX8/1491Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAkas Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description7849Gene ID7849Gene SymbolPAX8Specificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular carinomas, and atypical thyroid adenomas. PAX-8 is expressed in the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian	, , , , , , , , , , , , , , , , , , ,	•
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePAX8/1491Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid folicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid folicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid ovarian surface epithelial cells. PAX-8 is expressed in the thyroid (and associated carcinomas, put not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, put not normal ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.	Product Information	
services.         Storage       Store at 4C in the dark.         Clonality       Monoclonal         Clone       PAX8/1491         Preservative       0.05% Sodium Azide         Isotype       IgG2b Kappa         Conjugate       Alexa Fluor 647         Purity       Protein A or G purified         Buffer       50mM Sodium Borate         Product Description       Mouse         Gene ID       7849         Gene Symbol       PAX8         Species       Human         Marker       Renal Cell Marker         Specificity/Sensitivity       Recognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid systemes is, thyroid follicular cell development and expression of thyroid systemes is, thyroid follicular cell development and expression of thyroid systemes is, thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but on tormain surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometricid, and clear cell carcinomas, PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, A expension is reported in renal tubules	Unit Size	100 ul
ClonalityMonoclonalClonePAX8/1491Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerReal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), but on romal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubles as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Concentration	
ClonePAX8/1491Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerReal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular carinomas, and atypical thyroid agenesis, but or and simple ovarian incluse or ayrain and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thryroid folicular cell development and expression of thryroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid folicular cell development and expression of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, path a bit percentage of ovarian serous, endometrioid, and clear cell carcinomas, path a bit percentage of ovarian surface well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Clonality	Monoclonal
IsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of there of the factors and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, path expression is reported in renal tubules as well as renal cell carcinoma. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma. PAX-8 protein (exactImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Clone	PAX8/1491
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Isotype	IgG2b Kappa
Buffer       50mM Sodium Borate         Product Description       Host         Host       Mouse         Gene ID       7849         Gene Symbol       PAX8         Species       Human         Marker       Renal Cell Marker         Specificity/Sensitivity       Recognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.         Immunogen       Recombinant fragment (around aa 60-261) of human PAX8 protein (exact	Conjugate	Alexa Fluor 647
Product DescriptionHostMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas, non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Purity	Protein A or G purified
HostMouseGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Buffer	50mM Sodium Borate
Gene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Product Description	
Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinoma. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Host	Mouse
SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Gene ID	7849
MarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Gene Symbol	PAX8
Specificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenRecombinant fragment (around aa 60-261) of human PAX8 protein (exact	Species	Human
<ul> <li>box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.</li> <li>Immunogen</li> </ul>	Marker	Renal Cell Marker
	Specificity/Sensitivity	box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial
	Immunogen	



Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Protein Array
<b>Recommended Dilutions</b>	Western Blot, Immunohistochemistry-Paraffin, Protein Array
Application Notes	Optimal dilution of this antibody should be experimentally determined.





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 novus@novusbio.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: technical@novusbio.com Orders: orders@novusbio.com General: novus@novusbio.com

#### Products Related to NBP2-54486AF647

NBP1-43317AF647	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor® 647]
NBP2-51907-0.05mg	Recombinant Human PAX8 His Protein
288-TPN-025/CF	Thrombopoietin/THPO [Unconjugated]
NBL1-14132	PAX8 Overexpression Lysate

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54486AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

