

Product Datasheet

5-MethylCytosine Antibody (33D3) - BSA Free NBP2-54609

Unit Size: 100 ug

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-54609

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-54609




NBP2-54609**5-MethylCytosine Antibody (33D3) - BSA Free**

Product Information	
Unit Size	100 ug
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	33D3
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein A purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse 5-MethylCytosine Antibody (33D3) - BSA Free (NBP2-54609) is a monoclonal antibody validated for use in ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Species	Human, Mouse, Rat, Bovine
Immunogen	5-methylcytosine
Product Application Details	
Applications	Dot Blot, Immunocytochemistry/ Immunofluorescence, Methylated DNA Immunoprecipitation, Immunofluorescence, Surface Plasmon Resonance
Recommended Dilutions	Immunocytochemistry/ Immunofluorescence 1:500, Dot Blot 1:100, Methylated DNA Immunoprecipitation 0.5 - 1.0 ug/IP, Surface Plasmon Resonance, Immunofluorescence 1:500




Images


Immunocytochemistry/Immunofluorescence: 5-MethylCytosine Antibody (33D3) [NBP2-54609] - HeLa cells were stained with the antibody against 5-mC and with DAPI. Cells were fixed with 4% formaldehyde for 10 minutes and blocked with PBS/TX-100 containing 1% BSA. The cells were immunofluorescently labelled with the 5-mC antibody (middle) diluted 1:500 in blocking solution followed by an anti-mouse antibody conjugated to Alexa Fluor 594. The left panel shows staining of the nuclei with DAPI. A merge of the two stainings is shown on the right.




Surface Plasmon Resonance: 5-MethylCytosine Antibody (33D3) [NBP2-54609] - A synthesized biotin-labeled 5-mC conjugate was immobilized on a sensorchip. Briefly, two flowcells were prepared by sequential injections of EDC/NHS, streptavidin, and ethanolamine. One of these flowcells served as negative control, while biotinylated 5-mC conjugate was injected in the other one, to get an immobilization level of 55 response units (RU). All SPR experiments were performed, using HBS-N buffer (10 mM HEPES, 150 mM NaCl, pH 7.4), at a flow rate of 5 uL/min. Interaction assays involved injections of 2 different dilutions of the 5-mC monoclonal antibody over the biotinylated 5-mC conjugate and negative control surfaces, followed by a 3 minute washing step with HBS-N buffer. At the end of each cycle, the streptavidin surface was regenerated by injection of 0.1M citric acid (pH 3). The value of the dissociation constant (kd) obtained by global fitting and 1:1 Langmuir model is 65 nM.



Dot Blot: 5-MethylCytosine Antibody (33D3) [NBP2-54609] - To demonstrate the specificity of the antibody against 5-mC, a Dot blot analysis was performed using hmC, mC and C controls. 100 to 4 ng (equivalent of 5 to 0.2 pmol of C-bases) of the controls were spotted on a membrane. The antibody was used at a dilution of 1:300. Figure shows a high specificity of the antibody for the methylated control.



Methylated DNA Immunoprecipitation: 5-MethylCytosine Antibody (33D3) [NBP2-54609] - Analysis was performed on 1 ug fragmented human genomic DNA using 0.2 ug of the monoclonal antibody against 5-mC. The fragmented DNA was spiked with controls (methylated DNA (meDNA) as a positive and unmethylated DNA (unDNA) as a negative control) prior to performing the IP. QPCR was performed with primer sets specific for the methylated and unmethylated DNA controls, and for a known methylated (TSH2B) and unmethylated (GAPDH) genomic region. The figure shows the recovery expressed as a percent of input (the relative amount of immunoprecipitated DNA compared to input DNA after qPCR analysis).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-54609

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NBP2-62131	5-MethylCytosine ELISA Kit (Colorimetric)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54609

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

