Product Datasheet

ABCA1 Antibody (1276B) - BSA Free NBP2-54792

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54792

Updated 7/28/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54792



NBP2-54792

ABCA1 Antibody (1276B) - BSA Free


ABOAT Attibody (1270b) BOAT Tee		
Product Information		
Unit Size	0.1 mg	
Concentration	1.0 mg/ml	
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Clonality	Monoclonal	
Clone	1276B	
Preservative	0.02% Sodium Azide	
Isotype	IgG	
Purity	Protein A or G purified	
Buffer	PBS	

Product Description	
Host	Rabbit
Gene ID	19
Gene Symbol	ABCA1
Species	Human, Mouse
Immunogen	A synthetic peptide made to an internal portion of the human ABCA1 Antibody (1276B) (between amino acids 1220-1280) [UniProt# O95477]


Product Application Details	
Applications	Western Blot, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 2 ug/ml, Immunohistochemistry 1:50 - 1:100, Immunocytochemistry/ Immunofluorescence 10-20 ug/ml, Immunohistochemistry-Paraffin 1:50 - 1:100, Flow (Intracellular) 1-2.5 ug/mL

Images


Western Blot: ABCA1 Antibody (1276B) [NBP2-54792] - Mouse Raw264.7 and human THP-1 cells were treated with and without the LXR agonist TO901317 (1 uM) for 24 hours. Total protein was then separated on a 12% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 0.5 ug/ml anti-ABCA1 in 5% non-fat milk in TBST and detected with an anti-rabbit HRP secondary antibody using chemiluminescence.




Immunohistochemistry-Paraffin: ABCA1 Antibody (1276B) [NBP2-54792] - Analysis of a FFPE tissue section of mouse liver using ABCA1 antibody (1276B) Lot ckjv02 at 1:100 dilution. The primary antibody bound to ABCA1 antigen in the tissue section was detected using a HRP labeled secondary antibody and DAB reagent. Nuclei of the cells were counterstained with hematoxylin. This ABCA1 antibody generated specific staining in the membranes of the hepatocytes and the Kupffer cells. As expected for ABCA1, the staining signal was stronger in the Kupffer cells which are the tissue macrophages in liver.




Immunocytochemistry/Immunofluorescence: ABCA1 Antibody (1276B) [NBP2-54792] - HepG2 cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X TBS + 0.5% Triton X-100. The cells were incubated with anti-ABCA1 at 10 ug/ml overnight at 4C and detected with an anti-rabbit DyLight 488 (Green) at a 1:500 dilution. Alpha tubulin (DM1A) NB100-690 was used as a co-stain at a 1:1000 dilution and detected with an anti-mouse DyLight 550 (Red) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




Flow (Intracellular): ABCA1 Antibody (1276B) [NBP2-54792] - An intracellular stain was performed on HepG2 Cells with ABCA1 (1276B) antibody NBP2-54792 (blue) and a matched isotype control MAB1050 (orange). Cells were fixed with 4% paraformaldehyde, following fixation, cells were permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1 ug/mL for 30 minutes at room temperature, followed by rabbit IgG APC-conjugated secondary antibody (F0111, R&D Systems).



Immunocytochemistry/Immunofluorescence: ABCA1 Antibody (1276B) [NBP2-54792] - HeLa cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X TBS + 0.5% Triton X-100. The cells were incubated with anti-ABCA1 at 10 ug/ml overnight at 4C and detected with an anti-rabbit DyLight 488 (Green) at a 1:500 dilution. Alpha tubulin (DM1A) NB100-690 was used as a co-stain at a 1:1000 dilution and detected with an anti-mouse DyLight 550 (Red) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.



Immunohistochemistry-Paraffin: ABCA1 Antibody (1276B) [NBP2-54792] - Analysis of a FFPE tissue section of mouse liver using ABCA1 antibody (1276B) Lot ckjv0116091 at 1:50 dilution. The primary antibody bound to ABCA1 antigen in the tissue section was detected using a HRP labeled secondary antibody and DAB reagent. Nuclei of the cells were counterstained with hematoxylin. This ABCA1 antibody generated specific staining in the membranes of the hepatocytes and the Kupffer cells. As expected for ABCA1, the staining signal was stronger in the Kupffer cells which are the tissue macrophages in liver.



Publications

Zhao W, Fan J, Kulic I et al. Axl receptor tyrosine kinase is a regulator of apolipoprotein E Mol Brain 2020-05-04 [PMID: 32366277] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-54792

NBP3-11828 THP-1 TO901317 Treated / Untreated Cell Lysate

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54792

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

