Product Datasheet

WC1(+) gamma delta T cell Antibody (CACTB21A) NBP2-61081

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-61081

Updated 4/21/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-61081



NBP2-61081

WC1(+) gamma delta T cell Antibody (CACTB21A)


| VVC1(+) gamma delta T cell Antibody (CACTB21A) | | |
|--|---|--|
| Product Information | | |
| Unit Size | 0.1 mg | |
| Concentration | This product is unpurified. The exact concentration of antibody is not quantifiable. | |
| Storage | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. | |
| Clonality | Monoclonal | |
| Clone | CACTB21A | |
| Preservative | 0.09% Sodium Azide | |
| Isotype | IgG1 | |
| Purity | Unpurified | |
| Buffer | Ascites | |
| Product Description | | |
| Description | This monoclonal antibody is made in ascites form using mice. The ascites is collected and centrifuged to remove cells and large particles. The ascitic fluid is | |

| Product Description | |
|---------------------|---|
| Description | This monoclonal antibody is made in ascites form using mice. The ascites is collected and centrifuged to remove cells and large particles. The ascitic fluid is purified further by filtration through sequential filters with the last filter at 2 um. The concentration of the monoclonal antibody present in the ascites is determined with commercial radial immunodiffusion kits (1mg/ml). |
| Host | Mouse |
| Species | Bovine |
| Immunogen | Bovine ConA stimulated mononuclear leukocytes |
| Notes | This monoclonal antibody is made in ascites form using mice. The ascites is collected and centrifuged to remove cells and large particles. The ascitic fluid is purified further by filtration through sequential filters with the last filter at 2 um. The concentration of the monoclonal antibody present in the ascites is determined with commercial radial immunodiffusion kits (1mg/ml). |

| Product Application Details | |
|-----------------------------|------------------------------|
| Applications | Flow Cytometry |
| Recommended Dilutions | Flow Cytometry 1:10 - 1:1000 |

Images

Flow Cytometry: WC1(+) gamma delta T cell Antibody (CACTB21A) [NBP2-61081] - Obtained with bovine leukocytes







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-61081

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-61081

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

