# **Product Datasheet**

# CD42b/GPIb alpha Antibody (AK2) [PE/Cy7] NB100-63790PECY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-63790PECY7

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-63790PECY7



# NB100-63790PECY7

CD42b/GPIb alpha Antibody (AK2) [PE/Cy7]

Stora at 4C in the dark. Do not freeze.  Clonality Monoclonal  Clone AK2  Preservative 0.05% Sodium Azide  Isotype IgG1  Conjugate PE/Cy7  Purity Protein G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 2811  Gene Symbol GP1BA  Species Human  Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details  Applications ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit	CD42b/GPIb alpha Antibody (A	K2) [PE/Gy7]
Concentration  Please see the vial label for concentration. If unlisted please contact technic services.  Storage  Store at 4C in the dark. Do not freeze.  Clonality  Monoclonal  Clone  AK2  Preservative  Jo.5% Sodium Azide  Isotype  IgG1  Conjugate  PE/Cy7  Purity  Protein G purified  Buffer  PBS  Product Description  Host  Mouse  Gene ID  2811  Gene Symbol  GP1BA  Species  Human  Reactivity Notes  Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity  Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen  Made to Human CD42b  Product Application Details  Applications  ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions  Application Notes  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please court echnical support team at technical@novusbio.com if you have any	Product Information	
Storage Store at 4C in the dark. Do not freeze.  Clonality Monoclonal  Clone AK2  Preservative 0.05% Sodium Azide  Isotype IgG1  Conjugate PE/Cy7  Purity Protein G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 2811  Gene Symbol GP1BA  Species Human  Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details  Applications ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions  Application Notes  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please oc our technical support team at technical@novusbio.com if you have any	Unit Size	0.1 ml
Clonality Monoclonal Clone AK2 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate PE/Cy7 Purity Protein G purified Buffer PBS  Product Description Host Mouse Gene ID 2811 Gene Symbol GP1BA Species Human Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579). Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details Applications ELISA, Flow Cytometry, Immunoprecipitation Recommended Dilutions Flow Cytometry, ELISA, Immunoprecipitation Application Notes Optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please cour technical support team at technical@novusbio.com if you have any	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Clone AK2  Preservative 0.05% Sodium Azide  Isotype IgG1  Conjugate PE/Cy7  Purity Protein G purified  Buffer PBS  Product Description  Host Mouse Gene ID 2811  Gene Symbol GP1BA  Species Human  Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone Ak2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details  Applications ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions Flow Cytometry, ELISA, Immunoprecipitation  Application Notes Optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please or our technical support team at technical@novusbio.com if you have any	Storage	Store at 4C in the dark. Do not freeze.
Preservative   19G1   Conjugate   PE/Cy7   Purity   Protein G purified   Buffer   PBS   Product Description   Host   Mouse   Gene ID   2811   Gene Symbol   GP1BA   Species   Human   Reactivity Notes   Use in Mouse reported in scientific literature (PMID:29437579). Specificity/Sensitivity   Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1). Immunogen   Made to Human CD42b   Product Application Details   Application   ELISA, Flow Cytometry, Immunoprecipitation   Recommended Dilutions   Flow Cytometry, ELISA, Immunoprecipitation   Application Notes   Optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please occour technical support team at technical@novusbio.com if you have any	Clonality	Monoclonal
Isotype	Clone	AK2
Conjugate PE/Cy7  Purity Protein G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 2811  Gene Symbol GP1BA  Species Human  Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details  Applications ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions Flow Cytometry, ELISA, Immunoprecipitation  Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any	Preservative	0.05% Sodium Azide
Purity Protein G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 2811  Gene Symbol GP1BA  Species Human  Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details  Applications ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions Flow Cytometry, ELISA, Immunoprecipitation  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please court technical support team at technical@novusbio.com if you have any	Isotype	IgG1
Product Description Host Mouse Gene ID 2811 Gene Symbol GP1BA Species Human Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579). Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1). Immunogen Made to Human CD42b  Product Application Details Applications ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions Flow Cytometry, ELISA, Immunoprecipitation  Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please or our technical support team at technical@novusbio.com if you have any	Conjugate	PE/Cy7
Product Description Host Mouse  Gene ID 2811  Gene Symbol GP1BA  Species Human  Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details  Applications ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions Flow Cytometry, ELISA, Immunoprecipitation  Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please cour technical support team at technical@novusbio.com if you have any	Purity	Protein G purified
Host Mouse  Gene ID 2811  Gene Symbol GP1BA  Species Human  Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details  Applications ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions Flow Cytometry, ELISA, Immunoprecipitation  Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please cour technical support team at technical@novusbio.com if you have any	Buffer	PBS
Gene Symbol Gene Symbol Gene Symbol Gene Symbol Gene Symbol GP1BA Species Human Reactivity Notes Use in Mouse reported in scientific literature (PMID:29437579). Specificity/Sensitivity Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen Made to Human CD42b  Product Application Details Applications ELISA, Flow Cytometry, Immunoprecipitation Recommended Dilutions Flow Cytometry, ELISA, Immunoprecipitation  Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please cour technical support team at technical@novusbio.com if you have any	<b>Product Description</b>	
Gene Symbol  Species  Human  Reactivity Notes  Use in Mouse reported in scientific literature (PMID:29437579).  Specificity/Sensitivity  Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen  Made to Human CD42b  Product Application Details  Applications  ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions  Flow Cytometry, ELISA, Immunoprecipitation  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fit and permeabilization steps in your flow protocol may be required. Please cour technical support team at technical@novusbio.com if you have any	Host	Mouse
Species	Gene ID	2811
Reactivity Notes  Use in Mouse reported in scientific literature (PMID:29437579).  Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen  Made to Human CD42b  Product Application Details  Applications  ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions  Flow Cytometry, ELISA, Immunoprecipitation  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please cour technical support team at technical@novusbio.com if you have any	Gene Symbol	GP1BA
Specificity/Sensitivity  Recognises the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen  Made to Human CD42b  Product Application Details  Applications  ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions  Flow Cytometry, ELISA, Immunoprecipitation  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please coour technical support team at technical@novusbio.com if you have any	Species	Human
glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets (1).  Immunogen  Made to Human CD42b  Product Application Details  Applications  ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions  Flow Cytometry, ELISA, Immunoprecipitation  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please coour technical support team at technical@novusbio.com if you have any	Reactivity Notes	Use in Mouse reported in scientific literature (PMID:29437579).
Product Application Details  Applications  ELISA, Flow Cytometry, Immunoprecipitation  Recommended Dilutions  Application Notes  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please coour technical support team at technical@novusbio.com if you have any	Specificity/Sensitivity	glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor
Applications  ELISA, Flow Cytometry, Immunoprecipitation  Flow Cytometry, ELISA, Immunoprecipitation  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please coour technical support team at technical@novusbio.com if you have any	Immunogen	Made to Human CD42b
Recommended Dilutions  Flow Cytometry, ELISA, Immunoprecipitation  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please coour technical support team at technical@novusbio.com if you have any	<b>Product Application Details</b>	
Application Notes  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any	Applications	ELISA, Flow Cytometry, Immunoprecipitation
optimal results using our Tandem dyes, please avoid prolonged exposure to or extreme temperature fluctuations. These can lead to irreversible degrada or decoupling. When staining intracellular targets, specific attention to the fix and permeabilization steps in your flow protocol may be required. Please co our technical support team at technical@novusbio.com if you have any	Recommended Dilutions	Flow Cytometry, ELISA, Immunoprecipitation
1	Application Notes	optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NB100-63790PECY7

NBP1-97005PECY7 Mouse IgG1 Isotype Control (MG1) [PE/Cy7]

NBP1-85749PEP CD42b/GPIb alpha Recombinant Protein Antigen

210-TA-005 TNF-alpha [Unconjugated]

4067-GP-050 CD42b/GPIb alpha [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-63790PECY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

