

Product Datasheet

HLA Class I Antibody (MEM-147) [PE/Cy7] NB500-411PECY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB500-411PECY7

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB500-411PECY7



NB500-411PECY7

HLA Class I Antibody (MEM-147) [PE/Cy7]

| Product Information | |
|-----------------------------|--|
| Unit Size | 0.1 ml |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at 4C in the dark. Do not freeze. |
| Clonality | Monoclonal |
| Clone | MEM-147 |
| Preservative | 0.05% Sodium Azide |
| Isotype | IgG1 |
| Conjugate | PE/Cy7 |
| Purity | Protein A purified |
| Buffer | PBS |
| Product Description | |
| Host | Mouse |
| Gene ID | 3105 |
| Gene Symbol | HLA-A |
| Species | Human |
| Specificity/Sensitivity | This antibody (clone MEM-147) reacts with all human classical MHC Class I molecules in native cell-surface forms (e.g. it recognizes native HLA-A2 in cytofluorometry and immunoprecipitation but not in Western blotting). MHC Class I molecules (MHC Class Ia) are expressed on the surface of all human nucleated cell types. The antibody MEM-147 is positive in Western blotting (non-reducing conditions) only with most HLA-B and HLA-C molecules, but not HLA-A. Reactivity is very similar to the classical antibody W6/32. |
| Immunogen | PHA-activated peripheral blood lymphocytes |
| Product Application Details | |
| Applications | Flow Cytometry |
| Recommended Dilutions | Flow Cytometry |
| Application Notes | Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any questions. |





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB500-411PECY7

| | |
|-----------------|---|
| NBP1-97005PECY7 | Mouse IgG1 Isotype Control (MG1) [PE/Cy7] |
| 210-TA-005 | TNF-alpha [Unconjugated] |
| 285-IF-100 | IFN-gamma [Unconjugated] |
| 202-IL-010 | IL-2 [Unconjugated] |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB500-411PECY7

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

