Product Datasheet

Melan-A/MART-1 Antibody (A103 + M2-7C10 + M2-9E3) [PE/Cy5.5] NBP2-34546PECY55

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34546PECY55

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34546PECY55



NBP2-34546PECY55

Melan-A/MART-1 Antibody (A103 + M2-7C10 + M2-9E3) [PE/Cy5.5]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone A103 + M2-7C10 + M2-9E3 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG2b Lambda/IgG2b Kappa Conjugate PE/Cy5.5 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologous T lymphocytes. Seven other melanoma associated antigens recognize utologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subceillular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antib labels melanomas and other tumors showing melanocytic differentiation of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655) Product Application Details	
Concentration Please see the vial label for concentration. If unlisted please contact services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone A103 + M2-7C10 + M2-9E3 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG2b Lambda/IgG2b Kappa Conjugate PE/Cy5.5 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antible labels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone A103 + M2-7C10 + M2-9E3 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG2b Lambda/IgG2b Kappa Conjugate PE/Cy5.5 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as ((Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou T lymphocytes. Seven other melanoma associated antigens recognized to guatologougous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibo labels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain to fepithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
Clonality Monoclonal Clone A103 + M2-7C10 + M2-9E3 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG2b Lambda/IgG2b Kappa Conjugate PE/Cy5.5 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou T lymphocytes. Seven other melanoma associated antigens recognized sand other tumors showing melanocytic differentiat in melanosomes and endoplasmic reticulum. This monoclonal antibol labels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	echnical
Clone A103 + M2-7C10 + M2-9E3 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG2b Lambda/IgG2b Kappa Conjugate PE/Cy5.5 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou T lymphocytes. Seven other melanoma associated antigens recognize and tologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibol labels melanomas and other tumors showing melanocytic differentiation of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
Preservative Isotype IgG1 Kappa/IgG2b Lambda/IgG2b Kappa Conjugate PE/Cy5.5 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou. T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibo labels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Product Application Details	
Isotype IgG1 Kappa/IgG2b Lambda/IgG2b Kappa	
Conjugate PE/Cy5.5 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibolabels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Product Application Details	
Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologous T lymphocytes. Seven other melanoma associated antigens recognized autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibol labels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
Product Description Host Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibo labels melanomas and other tumors showing melanocytic differentiation also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Product Application Details	
Product Description Host Mouse Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou. T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibo labels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain to fepithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
Host Gene ID 2315 Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologous T lymphocytes. Seven other melanoma associated antigens recognizate autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, good BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antiboto labels melanomas and other tumors showing melanocytic differentiation also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655))	
Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologou T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibolabels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
Gene Symbol MLANA Species Human, Mouse, Rat, Canine Marker Melanoma Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologous T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibo labels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655))	
Species Human, Mouse, Rat, Canine Marker Melanoma Marker This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologous T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibolabels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
Marker Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologous T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibolabels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
Specificity/Sensitivity This antibody recognizes a protein doublet of 20-22kDa, identified as (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologous T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibol labels melanomas and other tumors showing melanocytic differentiat also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655))	
(Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is identified melanocyte differentiation antigen recognized by autologous T lymphocytes. Seven other melanoma associated antigens recognize autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, g BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 in melanosomes and endoplasmic reticulum. This monoclonal antibolabels melanomas and other tumors showing melanocytic differentiate also a useful positive-marker for angiomyolipomas. It does not stain to of epithelial, lymphoid, glial, or mesenchymal origin. Immunogen Recombinant full-length hMelan-A/MART-1 protein (A103; M2-7C10; (Uniprot: Q16655)	
(Uniprot: Q16655) Product Application Details	a newly s cytotoxic ed by p100, gp75, is present dy cocktail on. It is
• •	M2-9E3)
A P 4	
Applications Western Blot, Flow Cytometry, Flow (Intracellular), Immunocytochem Immunofluorescence, Immunohistochemistry, Immunohistochemistry CyTOF-ready	
Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytoc Immunofluorescence, Immunohistochemistry-Paraffin, Flow (Intracell CyTOF-ready	
Application Notes Optimal dilution of this antibody should be experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally optimal results using our Tandem dyes, please avoid prolonged experimentally optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined optimal results using our Tandem dyes, please avoid prolonged experimentally determined or extreme dyes, plea	sure to light egradation the fixation ase contact





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34546PECY55

NBP2-23223 Recombinant Human Melan-A/MART-1 His Protein

285-IF-100 IFN-gamma [Unconjugated]

NBP2-33535PEP Melan-A/MART-1 Recombinant Protein Antigen

202-IL-010 IL-2 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34546PECY55

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

