

Product Datasheet

Siglec-3/CD33 Antibody (C33/68) [PE/Cy5.5] NBP2-34579PECY55

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-34579PECY55

Updated 9/10/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-34579PECY55



NBP2-34579PECY55

Siglec-3/CD33 Antibody (C33/68) [PE/Cy5.5]

Product Information

| | |
|----------------------|---|
| Unit Size | 0.1 ml |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at 4C in the dark. Do not freeze. |
| Clonality | Monoclonal |
| Clone | C33/68 |
| Preservative | 0.05% Sodium Azide |
| Isotype | IgG1 Kappa |
| Conjugate | PE/Cy5.5 |
| Purity | Protein A purified |
| Buffer | PBS |

Product Description

| | |
|--------------------------------|---|
| Host | Mouse |
| Gene ID | 945 |
| Gene Symbol | CD33 |
| Species | Human |
| Specificity/Sensitivity | Recognizes a 67kDa glycoprotein, which is identified as CD33. It is a transmembrane protein of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. It belongs to the immunoreceptor tyrosine-based inhibitory motif (ITIM)-containing molecules able of recruiting protein tyrosine phosphatases SHP-1 and SHP-2 to signal assemblies; these ITIMs are also used for ubiquitin-mediated removal of the receptor from the cell surface. CD33 is expressed on cells of myelomonocytic lineage, binds sialic acid residues in N- and O-glycans on cell surfaces, and is a therapeutic target for acute myeloid leukemia. CD33 is expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. It is absent on platelets, lymphocytes, erythrocytes and hematopoietic stem cells. |
| Immunogen | Recombinant human CD33 protein |

Product Application Details

| | |
|------------------------------|--|
| Applications | Flow Cytometry |
| Recommended Dilutions | Flow Cytometry |
| Application Notes | Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any questions. |



Novus Biologicals USA

10730 E. Briarwood Avenue

Centennial, CO 80112

USA

Phone: 303.730.1950

Toll Free: 1.888.506.6887

Fax: 303.730.1966

novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave

Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400

Toll Free: 855.668.8722

Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane

Abingdon Science Park

Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15

Fax: (44) (0) 1235 533420

info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis.

Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34579PECY55

Earn gift cards/discounts by submitting a publication using this product:

www.novusbio.com/publications