# **Product Datasheet**

# beta 2-Microglobulin Antibody (246-E9.E7 (same as HLA.ABC.m2)) [PE/Cy7] NBP2-47700PECY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47700PECY7

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-47700PECY7



# NBP2-47700PECY7

beta 2-Microglobulin Antibody (246-E9.E7 (same as HLA.ABC.m2)) [PE/Cy7]

I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen  Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Applications  Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions  Flow Cytometry, Immunofluorescence, CyTOF-ready  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to ligh or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation	beta 2-Microglobulin Antibody (246-E9.E7 (same as HLA.ABC.m2)) [PE/Cy7]		
Please see the vial label for concentration. If unlisted please contact technical services.	Product Information		
Storage Store at 4C in the dark. Do not freeze.  Clonality Monoclonal  Clone 246-E9.E7 (same as HLA.ABC.m2)  Preservative 0.05% Sodium Azide  Isotype IgG2a Kappa  Conjugate PE/Cy7  Purity Protein A or G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 567  Gene Symbol BZM  Species Human, Primate (Negative)  Reactivity Notes Does not react with Non-Human primates.  Marker Renal Failure & Tumor Marker  Specificity/Sensitivity Recognizes a protein of 12kDa, identified as beta-microglobulin. Beta-microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigen complex. Human beta-2 microglobulin associated with HLA Clas I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Applications Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions Flow Cytometry, Immunofluorescence, CyTOF-ready Optimal results using our Tandem dyes, please avoid prolonged exposure to tig or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixaltian and permeabilization steps in your flow protocol may be required. Please conta our technic	Unit Size	0.1 ml	
Clone 246-E9.E7 (same as HLA.ABC.m2)  Preservative 0.05% Sodium Azide  Isotype IgG2a Kappa  Conjugate PE/Cy7  Purity Protein A or G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 567  Gene Symbol B2M  Species Human, Primate (Negative)  Reactivity Notes Does not react with Non-Human primates.  Marker Renal Failure & Tumor Marker  Specificity/Sensitivity  Recognizes a protein of 12kDa, identified as beta-microglobulin. Beta-microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. Petection of beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions Flow Cytometry, Immunofluorescence, CyTOF-ready  Application Notes  Product Application Details  Application Notes  Product Application Sepsion or Tandem dyes, please avoid prolonged exposure to ligit or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatic and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Concentration	· ·	
Clone 246-E9.E7 (same as HLA.ABC.m2)  Preservative 0.05% Sodium Azide  Isotype IgG2a Kappa  Conjugate PE/Cy7  Purity Protein A or G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 567  Gene Symbol B2M  Species Human, Primate (Negative)  Reactivity Notes Does not react with Non-Human primates.  Marker Renal Failure & Tumor Marker  Specificity/Sensitivity Recognizes a protein of 12kDa, identified as beta-microglobulin. Beta-microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Application Notes Plow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions Flow Cytometry, Immunofluorescence, CyTOF-ready  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to ligit or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatic and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Storage	Store at 4C in the dark. Do not freeze.	
Preservative   IgG2a Kappa	Clonality	Monoclonal	
IgG2a Kappa	Clone	246-E9.E7 (same as HLA.ABC.m2)	
Conjugate PE/Cy7  Purity Protein A or G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 567  Gene Symbol B2M  Species Human, Primate (Negative)  Reactivity Notes Does not react with Non-Human primates.  Marker Renal Failure & Tumor Marker  Specificity/Sensitivity Recognizes a protein of 12kDa, identified as beta-microglobulin. Beta-microglobulin non-covalently associates with the 44kDa chain to form the HLA Clast antigen complex. Human beta-2 microglobulin associated with HLA Clast I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin and does not react with nor-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Applications Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions Flow Cytometry, Immunofluorescence, CyTOF-ready  Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatic and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical @novusbio.com if you have any	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified  Buffer PBS  Product Description  Host Mouse  Gene ID 567  Gene Symbol B2M  Species Human, Primate (Negative)  Reactivity Notes Does not react with Non-Human primates.  Marker Renal Failure & Tumor Marker  Specificity/Sensitivity Recognizes a protein of 12kDa, identified as beta-microglobulin. Beta-microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, plateits, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to huma beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Applications Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions Flow Cytometry, Immunofluorescence, CyTOF-ready  Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to ligh or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatic and permeabilization steps in your flow protocol may be required. Please contain our technical with the protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Isotype	IgG2a Kappa	
PBS	Conjugate	PE/Cy7	
Product Description  Host Mouse  Gene ID 567  Gene Symbol B2M  Species Human, Primate (Negative)  Reactivity Notes Does not react with Non-Human primates.  Marker Renal Failure & Tumor Marker  Specificity/Sensitivity Recognizes a protein of 12kDa, identified as beta-microglobulin. Beta-microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigen complex. Human beta-2 microglobulin associated with HLA Clas I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Applications Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions Flow Cytometry, Immunofluorescence, CyTOF-ready  Application Notes Optimal results using our Tandem dyes, please avoid prolonged exposure to ligh or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatic and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Purity	Protein A or G purified	
Host   Mouse	Buffer	PBS	
Gene Symbol  Species  Human, Primate (Negative)  Reactivity Notes  Does not react with Non-Human primates.  Marker  Renal Failure & Tumor Marker  Specificity/Sensitivity  Recognizes a protein of 12kDa, identified as beta-microglobulin. Beta-microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigen complex. Human beta-2 microglobulin associated with HLA Clas I antigen sis expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen  Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Applications  Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions  Flow Cytometry, Immunofluorescence, CyTOF-ready  Application Notes  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to ligh or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatic and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Product Description		
Species	Host	Mouse	
Reactivity Notes   Does not react with Non-Human primates.	Gene ID	567	
Reactivity Notes   Does not react with Non-Human primates.	Gene Symbol	B2M	
Renal Failure & Tumor Marker	Species	Human, Primate (Negative)	
Recognizes a protein of 12kDa, identified as beta-microglobulin. Beta-microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigen complex. Human beta-2 microglobulin associated with HLA Clas I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen  Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Applications  Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions  Flow Cytometry, Immunofluorescence, CyTOF-ready  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to ligh or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatio and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Reactivity Notes	Does not react with Non-Human primates.	
microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigen complex. Human beta-2 microglobulin associated with HLA Clast I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitorin patients with HIV infection.  Immunogen  Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient  Product Application Details  Applications  Flow Cytometry, CyTOF-ready, Immunofluorescence  Recommended Dilutions  Flow Cytometry, Immunofluorescence, CyTOF-ready  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to ligh or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatic and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Marker	Renal Failure & Tumor Marker	
Product Application Details  Applications  Flow Cytometry, CyTOF-ready, Immunofluorescence  Flow Cytometry, Immunofluorescence, CyTOF-ready  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Specificity/Sensitivity	microglobulin non-covalently associates with the 44kDa chain to form the HLA Class I antigen complex. Human beta-2 microglobulin associated with HLA Class I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes. This monoclonal antibody is specific to human beta-2 microglobulin and does not react with non-human primate cells. This antibody reacts with all cell types excluding erythrocytes. Detection of beta-2 microglobulin in body fluids has been used as a tumor marker and for monitoring	
Applications  Flow Cytometry, CyTOF-ready, Immunofluorescence  Flow Cytometry, Immunofluorescence, CyTOF-ready  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Immunogen	Human PBL s from a T-cell acute lymphoblastic leukemia (T-ALL) patient	
Recommended Dilutions  Flow Cytometry, Immunofluorescence, CyTOF-ready  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contain our technical support team at technical@novusbio.com if you have any	Product Application Details		
Application Notes  Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Applications	Flow Cytometry, CyTOF-ready, Immunofluorescence	
optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Recommended Dilutions	Flow Cytometry, Immunofluorescence, CyTOF-ready	
	Application Notes	optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP2-47700PECY7

NBP1-96981PECY7 Mouse IgG2a Kappa Isotype Control (M2AK) [PE/Cy7]
NBP1-50840-0.01mg Recombinant Human beta 2-Microglobulin His Protein

210-TA-005 TNF-alpha [Unconjugated] KGE019 beta 2-Microglobulin [Biotin]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47700PECY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

