Product Datasheet

ECM-1/Secretory Component P85 Antibody (ECM1/792) [Alexa Fluor® 350] NBP2-47804AF350

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47804AF350

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-47804AF350



NBP2-47804AF350

ECM-1/Secretory Component P85 Antibody (ECM1/792) [Alexa Fluor® 350]

Unit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneECM1/792Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBufferSomM Sodium BorateProduct DescriptionKouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelial cells and epithelial differential differential secretory component and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differential and epithelial differential secretory component and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differential secretory component and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differential secretory component and the origen studying mucesal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneECM1/792Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1893HostMouseGene ID1893Gene SymbolECM1Specificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is a spell research tool for studying the distribution and level of both free and bound secretory component. Secretory component secretory component secretory component antibody is a spell research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component and tumorigenesis, all processes associated with differential secretory component and tumorigenesis, all processes associated with differential secretory component and tumorigenesis, all processes associated with differential secretory component and tumorigenesis, all processes associated with differential secretory component and tumorigenesis, all processes associated with differential secretory component at the secretory component and tumorigenesis, all processes associated with differential secretory component and tumorigenesis and processes associated with differential secretory component and tumorigenesis and processes associated with differential secretory component and tumorigenesis and processes associated with differe	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone ECM1/792 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 1893 Gene Symbol ECM1 Species Human, Rat Specificity/Sensitivity This monoclonal antibody reacts with a reduction-resistant epitope present in both free and SlgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component setpithelial differentiation. The Secretory component a	Unit Size	0.1 ml
ClonalityMonoclonalCloneECM1/792Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1893HostMouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processe associated with differential secretory component actibody is a sequil research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component met expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Concentration	
CloneECM1/792Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelian cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostHostMouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelial and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Clone	ECM1/792
ConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and turmorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelian, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Conjugate	Alexa Fluor 350
Product Description Host Mouse Gene ID 1893 Gene Symbol ECM1 Species Human, Rat Specificity/Sensitivity This monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression. Immunogen Recombinant full-length human ECM-1/Secretory Component P85 protein	Purity	Protein A or G purified
HostMouseGene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelium, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Buffer	50mM Sodium Borate
Gene ID1893Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelian, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Product Description	
Gene SymbolECM1SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Host	Mouse
SpeciesHuman, RatSpecificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelium, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Gene ID	1893
Specificity/SensitivityThis monoclonal antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Gene Symbol	ECM1
both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelium, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.ImmunogenRecombinant full-length human ECM-1/Secretory Component P85 protein	Species	Human, Rat
	Specificity/Sensitivity	both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelium, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component
	Immunogen	



N	otes	

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details

••	
Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Images

ECM-1/Secretory Component P85 Antibody (ECM1/792) [Alexa Fluor® 350] [NBP2-47804AF350] - Vial of Alexa Fluor 350 conjugated antibody. Alexa Fluor 350 is optimally excited at 346 nm by the UV laser (350 or 355 nm) and has an emission maximum of 442 nm.

Alexa Fluo	or® 350	
LASER (nm)	CII TEP	1
UV (350)	450/45	
		4)
EXCITATION MAX (nm)	EMISSION MAX (nm)	
	LASER (nm) UV (350)	UV (350) 450/45





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-47804AF350

NBP1-87449PEP	ECM-1/Secretory Component P85 Recombinant Protein Antigen
DMP900	MMP-9 [HRP]
3937-EC-050	ECM-1/Secretory Component P85 [Unconjugated]
AF808	Osteopontin/OPN Antibody [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47804AF350

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

