Product Datasheet

NGFR/TNFRSF16/p75NTR Antibody (NGFR5 + NTR/912) [Alexa Fluor® 594] NBP2-47967AF594

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47967AF594

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-47967AF594



NBP2-47967AF594

NGFR/TNFRSF16/p75NTR Antibody (NGFR5 + NTR/912) [Alexa Fluor® 594]

| Product Information 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone NGFR5+ NTR/912 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG1 Kappa Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotrophanal melanomas. NGFR is expressed in mature non-neuros and neurotrophoral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of bre | NGFR/TNFRSF10/p75NTR Altibody (NGFR5 + NTR/912) [Alexa Fluot® 594] | |
|--|--|---|
| Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone NGFR5 + NTR/912 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG1 Kappa Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal celd, and mycepithelial cells. Anti-NGFR stains the mycepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen | Product Information | |
| Storage Store at 4C in the dark. Clonality Monoclonal Clone NGFR5 + NTR/912 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG1 Kappa Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Unit Size | 0.1 ml |
| Clone NGFR5 + NTR/912 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG1 Kappa Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Concentration | ' |
| Clone NGFR5 + NTR/912 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa/IgG1 Kappa Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, phecofromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Storage | Store at 4C in the dark. |
| Preservative Isotype IgG1 Kappa/IgG1 Kappa Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Clonality | Monoclonal |
| IgG1 Kappa/IgG1 Kappa | Clone | NGFR5 + NTR/912 |
| Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Preservative | 0.05% Sodium Azide |
| Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Isotype | IgG1 Kappa/IgG1 Kappa |
| Buffer 50mM Sodium Borate | Conjugate | Alexa Fluor 594 |
| Product Description Host Mouse Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Purity | Protein A or G purified |
| Host Gene ID 4804 Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Buffer | 50mM Sodium Borate |
| Gene Symbol Reactivity Notes Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Product Description | |
| Gene Symbol NGFR Species Human, Primate, Mouse (Negative), Rat (Negative) Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Marker Soft Tissue Tumor Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Host | Mouse |
| Species | Gene ID | 4804 |
| Reactivity Notes Non-Human primates. Does not react with Mouse or Rat. Soft Tissue Tumor Marker It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Gene Symbol | NGFR |
| Marker Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti- NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Species | Human, Primate, Mouse (Negative), Rat (Negative) |
| Specificity/Sensitivity It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Reactivity Notes | Non-Human primates. Does not react with Mouse or Rat. |
| Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. Immunogen NGFR from A875 melanoma cells (NGFR5); Recombinant human | Marker | Soft Tissue Tumor Marker |
| | | Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts. |
| | Immunogen | |



Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

| Product Application Details | |
|-----------------------------|---|
| Applications | Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence |
| Recommended Dilutions | Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready |
| Application Notes | Optimal dilution of this antibody should be experimentally determined. |





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-47967AF594

NBP2-52198-0.05mg Recombinant Human NGFR/TNFRSF16/p75NTR His Protein

210-TA-005 TNF-alpha [Unconjugated] 367-NR-050 NGFR/TNFRSF16/p75NTR

DBD00 BDNF [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47967AF594

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

