

# Product Datasheet

## Cytokeratin 15 Antibody (LHK15) [PE/Cy7] NBP2-54462PECY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-54462PECY7](http://www.novusbio.com/NBP2-54462PECY7)

Updated 8/21/2022 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-54462PECY7](http://www.novusbio.com/reviews/destination/NBP2-54462PECY7)



**NBP2-54462PECY7**

Cytokeratin 15 Antibody (LHK15) [PE/Cy7]

| Product Information |   |
|---------------------|---|
| Unit Size           | 0.1 ml  |
| Concentration       | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage             | Store at 4C in the dark. Do not freeze.   |
| Clonality           | Monoclonal  |
| Clone               | LHK15   |
| Preservative        | 0.05% Sodium Azide  |
| Isotype             | IgG2a Kappa   |
| Conjugate           | PE/Cy7  |
| Purity              | Protein A or G purified   |
| Buffer              | PBS   |

| Product Description |   |
|---------------------|---|
| Host                | Mouse   |
| Gene ID             | 3866  |
| Gene Symbol         | KRT15   |
| Species             | Human, Rat, Bovine  |
| Marker              | Esophageal Squamous Cell Carcinoma Marker   |
| Immunogen           | A 17 amino acids synthetic peptide from the C- terminal of Human Cytokeratin 15 (Uniprot: P19012) |

| Product Application Details |  |
|-----------------------------|--|
| Applications                | Flow Cytometry   |
| Recommended Dilutions       | Flow Cytometry   |
| Application Notes           | Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at <a href="mailto:technical@novusbio.com">technical@novusbio.com</a> if you have any questions. |





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA

Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
novus@novusbio.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada

Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: technical@novusbio.com  
Orders: orders@novusbio.com  
General: novus@novusbio.com

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-54462PECY7](http://www.novusbio.com/reviews/submit/NBP2-54462PECY7)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

