Product Datasheet

Chorionic Gonadotropin alpha Chain (hCG alpha) Antibody (LHa/756) [Alexa Fluor® 750] NBP2-54500AF750

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-54500AF750

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-54500AF750

NBP2-54500AF750

Chorionic Gonadotropin alpha Chain (hCG alpha) Antibody (LHa/756) [Alexa Fluor® 750]

Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone LHa/756 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 750 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, folicle-stimulating hormone (FSH), throid-stimulating hormone (FSH), throid-stimulating hormone (FSH), throid-stimulating hormone adds (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit of the same receptor, however, the hCG beta subunit of the same into acids in sequence as the beta subunit on this es anae amino acids in sequence as the beta subunit of the same protector, throwever, the hCB beta subunit of the same breceptor, however, the hCB beta subunit on this manoecide and the hormone (FSH), it is nucates as and the hormone is a prohypept	· · ·	
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone LHa/756 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 750 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits nat are connected by two bridges. The alpha subunits of LH FSH, TSH, and hCG are identical, and contain 92 amino acids. ILHB that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunits an additional 24 amino acids in sequence as the beta subunit on the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin of IdGe-stimulating hormone (CSH), its is molaces alpha and beta subunits of LH as alpha acids and the hormone (ICSH), its is unedecert funue is also bee called Interstrital CeIH = Stimulating	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone LHa/756 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 750 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and hCG are identical, and contain 92 amino acids. The beta subunit contains the are connected by two bridges. The alpha subunit so fLH, FSH, TSH, and hCG are identical, and contain 92 amino acids (LHB) that confers its specific biologica action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit contains the same amino acids in sequence with the composition of their sugar moieties. LH is a synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the composition of their sugar moietes. LH is a synthesized and secreted by gonadotrophs in the mate, where LH has also been called Interstitiat Cell-Stimulating Hormone (ICSH), it simulates Leydig cell production of testosterone. LH is a useful marker	Unit Size	0.1 ml
Clonality Monoclonal Clone LHa/756 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 750 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 1081 Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits to tat, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunit contains the same amino acids in sequence as the beta subunit ontains an additional 24 amino acids and the hormones differ in the composition of their sugar moietes.LH is synthesized and secreted by gonadotroph follce-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it simulates Leydig cell production of teststerone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.	Concentration	•
CloneLHa/756Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1081Gene SymbolCGASpeciesHumanSpecificity/SensitivityThis monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (NG). The product of 21 amino acids (LHB) that confers its specific ado their subunit of 121 amino acids in sequence as the beta subunit contains an additional 24 amino acids an the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotropin folice-stimulating hormone (ICSH), it simulates the same receptor; however, the hCG beta subunit contains an additional 24 amino acids an acute rise of LH levels triggers ovulation. In the mane, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of teststerone. LH is a useful marker in classification of pitulary gland. In concert with the other productive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of teststerone. LH is a useful marker in classification of pitulary gland. In concert with the other pitulary gianadotropin folice-stimulating hormone (ICSH), it stimulates Leydig cell production of teststerone. LH is a useful marker in classification of pitulary fiturors and the study of pitulary disease. <th>Storage</th> <th>Store at 4C in the dark.</th>	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 750 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, folicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (NCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunit contains its responsible for interaction with the LH receptor. This beta subunit ontains in semino acids in sequence as the hCG abust work. LH is a beta subunit or 121 amino acids in sequences differ in the composition of their sugar moieties. LH is synthesized and secreted by gonadotropin follicle-stimulating hormone (ICSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers volution. In the female, an acute rise of LH levels triggers volution. In the female, an acute rise of LH levels triggers volution. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it simulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary gland. In concert with the other pituliary gisea	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 750PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1081Gene SymbolCGASpeciesHumanSpecificity/SensitivityThis monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunit on the asterion to fail and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contains 02 amino acids. The beta subunit on the asterio toolain of acids in sequence as the beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary turors and the study of pituitary disease.ImmunogenRecombinant full-length Chorionic Gonadotropin alpha Chain (hCG alpha)	Clone	LHa/756
Conjugate Alexa Fluor 750 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunit to rotains the same amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit ontains the same amino acids and the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituiary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary turors and the study of pituitary disease.	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Host Mouse Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadtropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits or tare annico acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit of heir sugar moieties. LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.	Isotype	IgG1 Kappa
Buffer 50mM Sodium Borate Product Description Mouse Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same are receptor, however, the hCG beta subunit of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other priutary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.	Conjugate	Alexa Fluor 750
Product Description Host Mouse Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunit vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is incuctive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.	Purity	Protein A or G purified
HostMouseGene ID1081Gene SymbolCGASpeciesHumanSpecificity/SensitivityThis monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.ImmunogenRecombinant full-length Chorionic Gonadotropin alpha Chain (hCG alpha)	Buffer	50mM Sodium Borate
Gene ID 1081 Gene Symbol CGA Species Human Specificity/Sensitivity This monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.	Product Description	
Gene SymbolCGASpeciesHumanSpecificity/SensitivityThis monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunit contains the same amino acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit of the sugar moieties.LH is synthesized and secreted by gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.ImmunogenRecombinant full-length Chorionic Gonadotropin alpha Chain (hCG alpha)	Host	Mouse
SpeciesHumanSpecificity/SensitivityThis monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.ImmunogenRecombinant full-length Chorionic Gonadotropin alpha Chain (hCG alpha)	Gene ID	1081
Specificity/SensitivityThis monoclonal antibody reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.ImmunogenRecombinant full-length Chorionic Gonadotropin alpha Chain (hCG alpha)	Gene Symbol	CGA
 sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease. 	Species	Human
Immunogen Recombinant full-length Chorionic Gonadotropin alpha Chain (hCG alpha)	Specificity/Sensitivity	sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties.LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary
	Immunogen	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-54500AF750

IC002S	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 750]
H00001081-Q01-10ug	Recombinant Human Chorionic Gonadotropin alpha Chain (hCG alpha) GST (N-Term) Protein
H00005670-P01-10ug	Recombinant Human PSG2 GST (N-Term) Protein
NBP2-75262	Human Chorionic Gonadotropin alpha Chain (hCG alpha) ELISA Kit (Colorimetric)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54500AF750

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com



www.novusbio.com



technical@novusbio.com