Product Datasheet

C19orf48 Knockout 293T Cell Lysate NBP2-65484

Unit Size: 2 Vials

Store at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-65484

Updated 10/20/2019 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-65484



NBP2-65484

| C19orf48 Knockout 293T Cell Lysate | |
|------------------------------------|---|
| Product Information | |
| Unit Size | 2 Vials |
| Concentration | Lyoph |
| Storage | Store at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles. |
| Reconstitution Instructions | To use as WB negative control, spin down briefly and resuspend in 100 uL 1xSDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Boil the lysate for 3 - 5 minutes before loading it onto gel. |
| Buffer | 0.1 mg cell homogenate lyophilized in RIPA buffer made with double-knockout cell lines. |
| Product Description | |
| Gene ID | 84798 |
| Gene Symbol | C19ORF48 |
| Species | Human |
| Preparation Method | Knockout achieved by using CRISPR/Cas9,-5 bp deletion in exon4 and 1 bp insertion in exon4 and Insertion of the selection cassette in exon4 |
| Notes | Powered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies. |
| Lysate Type | Knockout 293T Cell |
| Lysate Tissue | Kidney |
| Lysate Life Stage | Embryonic |
| Product Application Details | |
| Applications | Western Blot |
| Recommended Dilutions | Western Blot |
| Application Notes | You will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be |

| Product Application Details | |
|-----------------------------|---|
| Applications | Western Blot |
| Recommended Dilutions | Western Blot |
| Application Notes | You will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be stable at -20 degrees C for 12 months. Minimize freeze-thaw cycles. |





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-65484

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

