## **Product Datasheet**

# ATP5G3 Knockout 293T Cell Lysate NBP2-65538

Unit Size: 2 Vials

Store at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-65538

Updated 10/20/2019 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-65538



## NBP2-65538

ATP5G3 Knockout 293T Cell Lysate

ATP5G3 Knockout 293T Cell Lysate	
Product Information	
Unit Size	2 Vials
Concentration	Lyoph
Storage	Store at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.
Reconstitution Instructions	To use as WB negative control, spin down briefly and resuspend in 100 uL 1xSDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Boil the lysate for 3 - 5 minutes before loading it onto gel.
Buffer	0.1 mg cell homogenate lyophilized in RIPA buffer made with double-knockout cell lines.
Product Description	
Gene ID	518
Gene Symbol	ATP5G3
Species	Human
Preparation Method	Knockout achieved by using CRISPR/Cas9,11 bp deletion in exon 3
Notes	Powered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.
Lysate Type	Knockout 293T Cell
Lysate Tissue	Kidney
Lysate Life Stage	Embryonic
<b>Product Application Details</b>	
Applications	Western Blot
Recommended Dilutions	Western Blot
Application Notes	You will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. I yeate can be diluted with 1X SDS sample buffer and will be

<b>Product Application Details</b>	
Applications	Western Blot
Recommended Dilutions	Western Blot
	You will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be stable at -20 degrees C for 12 months. Minimize freeze-thaw cycles.





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Ltd
19 Barton Lane
Abingdon Science Park

Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-65538

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

