Product Datasheet

FAM128B Knockout 293T Cell Lysate NBP2-65606

Unit Size: 2 Vials

Store at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-65606

Updated 10/20/2019 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-65606



NBP2-65606

FAM128B Knockout 293T Cell Lysate

Product Information Unit Size 2 Vials Concentration Lyoph Storage Store at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles. Reconstitution Instructions To use as WB negative control, spin down briefly and resuspend in 100 uL 1XSDS sample buffer (2% SDS, 60 mM Tris-HCI pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Boil the lysate for 3 - 5 minutes before loading it onto gel. Buffer 0.1 mg cell homogenate lyophilized in RIPA buffer made with double-knockout cell lines. Product Description Gene ID Gene Symbol MZT2B Species Human Preparation Method Knockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1 Notes Powered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies. Lysate Type Knockout 293T Cell Lysate Life Stage Embryonic Product Application Details Vou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of parental cell lysate. Lysate can be diuted with 1X SDS sample buffer and will be stable at -20 degrees C for 12 months. Minimize freeze-thaw cycles.	-	
ConcentrationLyophStorageStore at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.Reconstitution InstructionsTo use as WB negative control, spin down briefly and resuspend in 100 uL 1xSDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Boil the lysate for 3 - 5 	Product Information	
StorageStore at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.Reconstitution InstructionsStore at -20C short term. Aliquot and store at -80C long term. Avoid freeze-thaw cycles.Reconstitution InstructionsTo use as WB negative control, spin down briefly and resuspend in 100 uL 1xSDS sample buffer (2% SDS, 60 mM Tris-HCI pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Boil the lysate for 3 - 5 minutes before loading it onto gel.Buffer0.1 mg cell homogenate lyophilized in RIPA buffer made with double-knockout cell lines.Product Description80097Gene ID80097Gene SymbolMZT2BSpeciesHumanPreparation MethodKnockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TypeKidneyLysate Life StageEmbryonicProduct Application DetailsWestern Blot Recommended DilutionsApplication NotesWestern BlotApplication NotesYou will receive 1 vial (100ug) of fraerent acid lysate can be diluted with 1X SDS sample buffer and will be	Unit Size	2 Vials
cycles.Reconstitution InstructionsTo use as WB negative control, spin down briefly and resuspend in 100 uL 1xSDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Boil the lysate for 3 - 5 minutes before loading it onto gel.Buffer0.1 mg cell homogenate lyophilized in RIPA buffer made with double-knockout cell lines.Product Description80097Gene ID80097Gene symbolMZT2BSpeciesHumanPreparation MethodKnockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Concentration	Lyoph
1xSDS sample buffer (2% SDS, 60 mM Tris-HCI pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Boil the lysate for 3 - 5 minutes before loading it onto gel.Buffer0.1 mg cell homogenate lyophilized in RIPA buffer made with double-knockout cell lines.Product Description60097Gene ID80097Gene SymbolMZT2BSpeciesHumanPreparation MethodKnockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TipsueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplication NotesYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Storage	
cell lines.cell lines.Product DescriptionGene ID80097Gene SymbolMZT2BSpeciesHumanPreparation MethodKnockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TypeKnockout 293T CellLysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Reconstitution Instructions	1xSDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Boil the lysate for 3 - 5
Gene ID80097Gene SymbolMZT2BSpeciesHumanPreparation MethodKnockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TypeKnockout 293T CellLysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsVou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Buffer	0.1 mg cell homogenate lyophilized in RIPA buffer made with double-knockout cell lines.
Gene SymbolMZT2BSpeciesHumanPreparation MethodKnockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TypeKnockout 293T CellLysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsWestern BlotApplication NotesYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Product Description	
SpeciesHumanPreparation MethodKnockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TypeKnockout 293T CellLysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsVou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Gene ID	80097
Preparation MethodKnockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TypeKnockout 293T CellLysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Gene Symbol	MZT2B
NotesPowered by EDIGENE. Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TypeKnockout 293T CellLysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Species	Human
Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative control for western blots and to confirm the specificity of antibodies.Lysate TypeKnockout 293T CellLysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsWestern BlotApplication NotesYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Preparation Method	Knockout achieved by using CRISPR/Cas9,29 bp deletion in exon 1
Lysate TissueKidneyLysate Life StageEmbryonicProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsWestern BlotApplication NotesYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Notes	Validation of antibody specificity is critical and verification of antibody performance against knockout samples is one way to guarantee that an antibody recognizes a specific target. Novus' KO cell lysate can be used as a negative
Lysate Life StageEmbryonicProduct Application DetailsApplicationsWestern BlotRecommended DilutionsWestern BlotApplication NotesYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Lysate Type	Knockout 293T Cell
Product Application Details Applications Western Blot Recommended Dilutions Western Blot Application Notes You will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Lysate Tissue	Kidney
ApplicationsWestern BlotRecommended DilutionsWestern BlotApplication NotesYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Lysate Life Stage	Embryonic
Recommended Dilutions Western Blot Application Notes You will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Product Application Details	
Application NotesYou will receive 1 vial (100ug) of knockout cell lysate and 1 vial (100ug) of Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Applications	Western Blot
Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be	Recommended Dilutions	Western Blot
	Application Notes	Parental cell lysate. Lysate can be diluted with 1X SDS sample buffer and will be





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-65606

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

