

Product Datasheet

TNF RII/TNFRSF1B Antibody (JM113-01) NBP2-67218

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-67218

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-67218



NBP2-67218

TNF RII/TNFRSF1B Antibody (JM113-01)

Product Information

Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JM113-01
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol

Product Description


Description	Novus Biologicals Rabbit TNF RII/TNFRSF1B Antibody (JM113-01) (NBP2-67218) is a recombinant monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	7133
Gene Symbol	TNFRSF1B
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within Human TNF RII/TNFRSF1B aa 426-461 / 461. (SwissProt: P20333 Human; SwissProt: P25119 Mouse; SwissProt: Q80WY6 Rat)

Product Application Details


Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500-1:2000, Flow Cytometry 1:50-1:100, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:50-1:200, Immunoprecipitation, Immunohistochemistry-Paraffin 1:50-1:200, Immunohistochemistry-Frozen 1:200

Images


Western Blot: TNF RII/TNFRSF1B Antibody (JM113-01) [NBP2-67218] - Western blot analysis of TNF RII/TNFRSF1B on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:5,000 dilution was used for 1 hour at room temperature. Positive control: Lane 2: MCF-7 cell lysate Lane 2: Jurkat cell lysate Lane 2: Hela cell lysate




Immunocytochemistry/Immunofluorescence: TNF RII/TNFRSF1B Antibody (JM113-01) [NBP2-67218] - Staining TNF Receptor II in SW480 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: TNF RII/TNFRSF1B Antibody (JM113-01) [NBP2-67218] - Analysis of paraffin-embedded mouse kidney tissue anti-TNF Receptor II antibody. Counter stained with hematoxylin.




Flow Cytometry: TNF RII/TNFRSF1B Antibody (JM113-01) [NBP2-67218] - Analysis of HL-60 cells with TNF Receptor II antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.




Immunocytochemistry/Immunofluorescence: TNF RII/TNFRSF1B Antibody (JM113-01) [NBP2-67218] - Staining TNF Receptor II in Hela cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Immunocytochemistry/Immunofluorescence: TNF RII/TNFRSF1B Antibody (JM113-01) [NBP2-67218] - Staining TNF Receptor II in MCF-7 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Immunohistochemistry-Paraffin: TNF RII/TNFRSF1B Antibody (JM113-01) [NBP2-67218] - Analysis of paraffin-embedded human kidney tissue using anti-TNF Receptor II antibody. Counter stained with hematoxylin.



Immunohistochemistry-Paraffin: TNF RII/TNFRSF1B Antibody (JM113-01) [NBP2-67218] - Analysis of paraffin-embedded human uterus tissue using anti-TNF Receptor II antibody. Counter stained with hematoxylin.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-67218

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-35182-5ug	Recombinant Mouse TNF RII/TNFRSF1B Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-67218

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

