Product Datasheet

PUMA Antibody (SR42-09) NBP2-67326

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-67326

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-67326



NBP2-67326

PUMA Antibody (SR42-09)


,	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	SR42-09
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol

Product Description	
Description	Novus Biologicals Rabbit PUMA Antibody (SR42-09) (NBP2-67326) is a recombinant monoclonal antibody validated for use in IHC, WB, Flow and ICC/IF. Anti-PUMA Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	27113
Gene Symbol	BBC3
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within Human PUMA aa 153-193 / 193. (SwissProt: Q9BXH1 Human; SwissProt: Q99ML1 Mouse; SwissProt: Q80ZG6 Rat)

Product Application Details	
	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
	Western Blot 1:500-1:2000, Flow Cytometry 1:50-1:100, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:50-1:200, Immunohistochemistry-Paraffin 1:50-1:200


Images

Western Blot: PUMA Antibody (SR42-09) [NBP2-67326] - Analysis of PUMA on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:5,000 dilution was used for 1 hour at room temperature. Positive control: Lane 1: Hela cell lysate Lane 2: K562 cell lysate







Immunocytochemistry/Immunofluorescence: PUMA Antibody (SR42-09) [NBP2-67326] - Staining PUMA in SKOV-3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: PUMA Antibody (SR42-09) [NBP2-67326] - Immunohistochemical analysis of paraffin-embedded mouse small intestine tissue using anti-PUMA antibody. The section was pretreated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BS




Flow Cytometry: PUMA Antibody (SR42-09) [NBP2-67326] - Analysis of Hela cells with PUMA antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.




Immunocytochemistry/Immunofluorescence: PUMA Antibody (SR42-09) [NBP2-67326] - Staining PUMA in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: PUMA Antibody (SR42-09) [NBP2-67326] - Analysis of paraffin-embedded human breast carcinoma tissue using anti-PUMA antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: PUMA Antibody (SR42-09) [NBP2-67326] - Analysis of paraffin-embedded human gastric carcimnoma tissue using anti-PUMA antibody. Counter stained with hematoxylin.



Immunohistochemistry-Paraffin: PUMA Antibody (SR42-09) [NBP2-67326] - Analysis of paraffin-embedded mouse small intestine tissue using anti-PUMA antibody. Counter stained with hematoxylin.



Immunohistochemistry-Paraffin: PUMA Antibody (SR42-09) [NBP2-67326] - Analysis of paraffin-embedded mouse stomach tissue using anti-PUMA antibody. Counter stained with hematoxylin.



Publications

Reina-Ortiz C, Mozas MP, Ovelleiro D et al. Dynamic Changes in miRNA Expression during the Generation of Expanded and Activated NK Cells International journal of molecular sciences 2023-08-31 [PMID: 37686362] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-67326

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NB500-261PEP PUMA Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-67326

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

