

Product Datasheet

eIF2A [p Ser51] Antibody (SZ01-06) NBP2-67353

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-67353

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-67353



NBP2-67353

eIF2A [p Ser51] Antibody (SZ01-06)

Product Information

| | |
|----------------------|--|
| Unit Size | 100 ul |
| Concentration | 1 mg/ml |
| Storage | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. |
| Clonality | Monoclonal |
| Clone | SZ01-06 |
| Preservative | 0.05% Sodium Azide |
| Isotype | IgG |
| Purity | Protein A purified |
| Buffer | TBS (pH7.4), 0.05% BSA, 40% Glycerol |

Product Description


| | |
|--------------------|---|
| Host | Rabbit |
| Gene ID | 83939 |
| Gene Symbol | EIF2A |
| Species | Human, Mouse, Rat |
| Immunogen | Synthetic phospho-peptide corresponding to residues surrounding Ser51 of Human eIF2A. (SwissProt: P05198 Human; SwissProt: Q6ZWX6 Mouse; SwissProt: P68101 Rat) |

Product Application Details


| | |
|------------------------------|---|
| Applications | Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation |
| Recommended Dilutions | Western Blot 1:500-1:2000, Flow Cytometry 1:50-1:100, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:50-1:200, Immunoprecipitation, Immunohistochemistry-Paraffin 1:50-1:500 |

Images


Western Blot: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of eIF2A [p Ser51] on THP-1 cell lysates. Lane 1: THP-1 cells, whole cell lysate, 10ug/lane Lane 2: THP-1 cells treated with 2.8ug/ul lambda-PP for 30 minutes, whole cell lysates, 10ug/lane All lanes : Anti-eIF2A [p Ser51]antibody at 1/500 dilution. Anti-GAPDH antibody at 1/10,000 dilution. Goat Anti-Rabbit IgG H&L (HRP) at 1/200,000 dilution. Predicted band size: 36 kDa Observed band size: 36 kDa Blocking and diluting buffer: 5% BSA. Exposure time: 3 minutes 43 seconds




Immunocytochemistry/Immunofluorescence: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Staining of Hela cells (green) using NBP2-67353. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of paraffin-embedded human breast carcinoma tissue with Rabbit anti-eIF2A [p Ser51] antibody washed with ddH₂O and PBS, and then probed with the primary antibody at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Flow Cytometry: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of Hela cells with Phospho-eIF-2a(S51) antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.




Western Blot: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of Phospho-eIF-2a(S51) on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:5,000 dilution was used for 1 hour at room temperature. Positive control: Lane 1: HUVEC cell lysate Lane 2: Hela cell lysate Lane 3: PC-3M cell lysate




Immunohistochemistry-Paraffin: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of paraffin-embedded human liver tissue using anti-Phospho-eIF-2a(S51) antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of paraffin-embedded human pancreas tissue using anti-Phospho-eIF-2a(S51) antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of paraffin-embedded mouse brain tissue using anti-Phospho-eIF-2a(S51) antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of paraffin-embedded mouse pancreas tissue using anti-Phospho-eIF-2a(S51) antibody. Counter stained with hematoxylin.




Immunohistochemistry-Paraffin: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of paraffin-embedded mouse placenta tissue using anti-Phospho-eIF-2a(S51) antibody. Counter stained with hematoxylin.



Immunohistochemistry-Paraffin: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Immunohistochemical analysis of paraffin-embedded human prostate carcinoma tissue using anti-eIF2A antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (ET1603-14, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunohistochemistry-Paraffin: eIF2A [p Ser51] Antibody (SZ01-06) [NBP2-67353] - Analysis of paraffin-embedded mouse brain tissue with Rabbit anti-eIF2A [p Ser51] antibody washed with ddH2O and PBS, and then probed with the primary antibody at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Publications

Chen Y, Tristan CA, Lu C et al. A Versatile Polypharmacology Platform Promotes Cytoprotection and Viability of Human Pluripotent and Differentiated Cells Nat Methods 2021-05-04 [PMID: 33941937] (Simple Western)

Kornicka-Garbowska K, Bourebaba L, ROcken M, Marycz K Sex Hormone Binding Globulin (SHBG) Mitigates ER Stress in Hepatocytes In Vitro and Ex Vivo Cells 2021-03-30 [PMID: 33808055] (WB, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-67353

| | |
|---------------|---|
| HAF008 | Goat anti-Rabbit IgG Secondary Antibody [HRP] |
| NB7160 | Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP] |
| NBP2-24891 | Rabbit IgG Isotype Control |
| NBP1-84879PEP | eIF2A Recombinant Protein Antigen |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-67353

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

