

# Product Datasheet

## Rabbit anti-Duck IgA/IgG/IgM (H+L) Secondary Antibody NBP2-69291

Unit Size: 1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-69291](http://www.novusbio.com/NBP2-69291)

Updated 11/2/2024 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-69291](http://www.novusbio.com/reviews/destination/NBP2-69291)



**NBP2-69291****Rabbit anti-Duck IgA/IgG/IgM (H+L) Secondary Antibody**

| <b>Product Information</b>         |                                                                                                                                                |
|------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Unit Size</b>                   | 1 ml                                                                                                                                           |
| <b>Concentration</b>               | LYOPH mg/ml                                                                                                                                    |
| <b>Storage</b>                     | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.                                                         |
| <b>Clonality</b>                   | Polyclonal                                                                                                                                     |
| <b>Preservative</b>                | No Preservative                                                                                                                                |
| <b>Reconstitution Instructions</b> | Reconstitute by adding 1 ml sterile distilled water. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). |
| <b>Purity</b>                      | Whole Antiserum                                                                                                                                |
| <b>Buffer</b>                      | Stable whole antiserum                                                                                                                         |

| <b>Product Description</b>     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|--------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>             | The lyophilized antiserum is shipped at ambient temperature and may be stored at 4C; prolonged storage at or below -20C. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at 4C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product. |
| <b>Host</b>                    | Rabbit                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Species</b>                 | Duck                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Reactivity Notes</b>        | Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.                                                                                                                                                                                                                                                                             |
| <b>Specificity/Sensitivity</b> | The reactivity of the antiserum is directed to the Fc and Fab subunits of the major immunoglobulins. It does not react with any non-Ig protein in duck serum, as tested by immunoelectrophoresis and double radial immunodiffusion.                                                                                                                                                                                                                                                                       |
| <b>Immunogen</b>               | Purified normal Ig fractions isolated from pooled duck serum. Freund's complete adjuvant is used in the first step of the immunization procedure.                                                                                                                                                                                                                                                                                                                                                         |

| <b>Product Application Details</b> |                                                                                                                                                                                                                                                                                                                                |
|------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Applications</b>                | Immunoelectrophoresis                                                                                                                                                                                                                                                                                                          |
| <b>Recommended Dilutions</b>       | Immunoelectrophoresis                                                                                                                                                                                                                                                                                                          |
| <b>Application Notes</b>           | <p>Precipitation assays. In immunoelectrophoresis use 2 ul or equivalent against 120 ul antiserum. In double radial immunodiffusion use a rosette arrangement with 10 ul antiserum in a 3 mm diameter center well and 2 ul serum samples (neat and diluted) in 2 mm diameter peripheral wells.</p> <p>Precipitation assays</p> |





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

[www.novusbio.com](http://www.novusbio.com)  
Technical Support: [nb-technical@bio-techne.com](mailto:nb-technical@bio-techne.com)  
Orders: [nb-customerservice@bio-techne.com](mailto:nb-customerservice@bio-techne.com)  
General: [novus@novusbio.com](mailto:novus@novusbio.com)

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-69291](http://www.novusbio.com/reviews/submit/NBP2-69291)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

