Product Datasheet

Rabbit anti-Monkey IgG F(ab) Secondary Antibody NBP2-69443

Unit Size: 10 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-69443

Updated 11/4/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-69443



NBP2-69443

Rabbit anti-Monkey IgG F(ab) Secondary Antibody

10 mg
LYOPH mg/ml
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Polyclonal
No Preservative
Reconstitute by by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below - 20C.
Ion exchange chromatography
Lyophilized from PBS (pH 7.2)
The lyophilized IgG fraction is shipped at ambient temperature and may be stored at 4C; prolonged storage at or below -20C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at 4C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.
Rabbit
Monkey
Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail, however in immunoelectrophoresis a reaction with Human has been observed.
The antiserum reacts with the Fab subunit of intact IgG, IgA and IgM and other Ig classes of both light chain types, with their Fab or F(ab')2 subunits and with free
light chains of kappa and lambda type as tested in immunoelectrophoresis and double radial immunodiffusion. The presence of activity to the common Fab subunit may result in the staining of Ig bound to Fc-receptors on non-lymphoid cells.
light chains of kappa and lambda type as tested in immunoelectrophoresis and double radial immunodiffusion. The presence of activity to the common Fab subunit may result in the staining of Ig bound to Fc-receptors on non-lymphoid
light chains of kappa and lambda type as tested in immunoelectrophoresis and double radial immunodiffusion. The presence of activity to the common Fab subunit may result in the staining of Ig bound to Fc-receptors on non-lymphoid cells.Purified Fab from normal IgG isolated from pooled Rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization
light chains of kappa and lambda type as tested in immunoelectrophoresis and double radial immunodiffusion. The presence of activity to the common Fab subunit may result in the staining of Ig bound to Fc-receptors on non-lymphoid cells.Purified Fab from normal IgG isolated from pooled Rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization
light chains of kappa and lambda type as tested in immunoelectrophoresis and double radial immunodiffusion. The presence of activity to the common Fab subunit may result in the staining of Ig bound to Fc-receptors on non-lymphoid cells.Purified Fab from normal IgG isolated from pooled Rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.Western Blot, Dot Blot, ELISA, Immunoblotting, Immunocytochemistry/





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-69443

NBP2-62014

IgG F(ab) Antibody (RMG01)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-69443

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

