Product Datasheet

Goat anti-Chicken IgM Fc Secondary Antibody [HRP] NBP2-69675

Unit Size: 1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-69675

Updated 11/2/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-69675



NBP2-69675

Goat anti-Chicken IgM Fc Secondary Antibody [HRP]	
Product Information	
Unit Size	1 ml
Concentration	LYOPH mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	No Preservative
Reconstitution Instructions	Reconstitute by by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below - 20C.
Conjugate	HRP
Purity	Ion exchange chromatography
Buffer	Lyophilized from PBS (pH 7.2)
Product Description	
Description	The lyophilized conjugate is shipped at ambient temperature and may be stored at 4C; prolonged storage at or below -20C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at 4C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.
Host	Goat
Species	Chicken
Reactivity Notes	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this immunoconjugate has not been tested in detail.
Specificity/Sensitivity	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in chicken serum, as tested by immunoelectrophoresis and double radial immunodiffusion.
Immunogen	Purified IgM isolated from chicken serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Product Application Details	
Applications	Western Blot, Dot Blot, ELISA, Immunohistochemistry-Paraffin, Immunocytochemistry



Western Blot, ELISA, Immunohistochemistry-Paraffin, Dot Blot, Immunocytochemistry

Recommended Dilutions



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-69675

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

