Product Datasheet

MUC6 Antibody (IHC626) NBP2-75985-0.1ml

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-75985

Updated 7/22/2020 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-75985



NBP2-75985-0.1ml

MUC6 Antibody (IHC626)


- ,	
Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark. Do not freeze.
Clonality	Monoclonal
Clone	IHC626
Preservative	0.1% Sodium Azide
Isotype	IgG2a
Purity	Protein A purified
Buffer	Tris Buffer, pH 7.3 - 7.7, with 1% BSA
Target Molecular Weight	252 kDa
Product Description	
Host	Mouse

Host	Mouse	
Gene ID	4588	
Gene Symbol	MUC6	
Species	Human	
Immunogen	Recombinant Human MUC6	
Product Application Details		
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin	

Applications	Immunohistochemistry, Immunohistochemistry-Paraffin	
Recommended Dilutions	Immunohistochemistry 1:100 - 1:200, Immunohistochemistry-Paraffin 1:100 - 1:200	
Application Notes	Visualization of the antigen present in tissue sections is accomplished in a multi- step immunohistochemical staining process, in conjunction with a horseradish peroxidase (HRP) or alkaline phosphatase (AP) linked detection system. Positive Control: Stomach. The 7mL size is a pre-diluted size and no additional dilutions are required before using this item for the intended application.	

Images

Immunohistochemistry-Paraffin: MUC6 Antibody (IHC626) [NBP2-75985] - Staining of MUC6 (IHC626) in stomach.





Procedures

Immunohistochemistry protocol for MUC6 Antibody (NBP2-75985)

Immunohistochemistry Protocol for MUC6 Antibody (NBP2-75985): https://www.novusbio.com/products/muc6-antibody-ihc626_nbp2-75985

Immunohistochemistry Protocol

Specimen Collection and Preparation for Analysis

Each tissue section should be fixed with 10% neutral buffered formalin, cut to the applicable thickness (4um), and placed on a glass slide that is positively charged. The prepared slide may then be baked for a minimum of 30 minutes in a 53-65 degrees C oven (do not exceed 24 hours).

Recommended Staining Protocols

Manual Use:

- 1. Pretreatment: Perform heat-induced epitope retrieval (HIER) at pH 9 for 10 to 30 minutes.
- 2. Peroxide Block: Block in peroxidase blocking solution for 5 minutes at room temperature. (Not required if using Alkaline Phosphatase System.)
- 3. Primary Antibody: Apply antibody directly (Predilute) or dilute antibody at 1:100-1:200 (Concentrate) before applying. Incubate antibody for 10 to 30 minutes at room temperature.
- 4. Secondary Antibody: Incubate for 20 to 30 minutes at room temperature.
- 5. Substrate Development: Incubate DAB or Fast Red for 5 to 10 minutes at room temperature.
- 6. Counterstain: Counterstain with hematoxylin for 0.5 to 5 minutes, depending on the hematoxylin used. Rinse with distilled water and blueing solution for 30 seconds.
- 7. Dehydrate and apply coverslip.

Automated Staining System:

The stated primary antibody has been optimized and validated using the BOND-MAX fully automated IHC & ISH stainer manufactured by Leica Biosystems, applying IHC Protocol F. The following edits are recommended for the protocol:

- a) Marker Incubation Time: 30 minutes
- b) Heat-induced epitope retrieval (HIER) is recommended using Leica Bond ER Solution 2 for 30 minutes.
- c) Move Peroxide Block step to after Polymer and before Mixed DAB Refine. For all other automated IHC staining systems, refer to the corresponding user manual for specific instructions.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Error: Subreport could not be shown.

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-75985

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

