Product Datasheet

LHR Antibody NBP2-76804

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-76804

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-76804



NBP2-76804


LHR Antibody

LHR Antibody	
0.1 mg	
0.5 mg/ml	
Store at -20C. Avoid freeze-thaw cycles.	
Polyclonal	
0.02% Sodium Azide	
IgG	
Immunogen affinity purified	
Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA	
Product Description	
Novus Biologicals Goat LHR Antibody (NBP2-76804) is a polyclonal antibody validated for use in WB, ELISA, Flow and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.	
Goat	
3973	
LHCGR	
Human	
Peptide with sequence CQGTALLDKTRYTE corresponding to C-Terminus according to NP_000224.2.	
Product Application Details	
Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Peptide ELISA	
Western Blot 0.3 - 0.1 ug/ml, Flow Cytometry 10 ug/ml, Immunocytochemistry/ Immunofluorescence, Peptide ELISA Detection Limit 1:32000	
WB: Preliminary experiments gave an approx. 35 kDa band in lysates of human brain, placenta and prostate and of cell lines Daudi, HeLa and K562 after 0.1 ug/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 78.6 kDa band according to NP_000224.2. The 35 kDa band was successfully blocked by incubation with the immunizing peptide.	




Images

Flow Cytometry: LHR Antibody [NBP2-76804] - Analysis of paraformaldehyde fixed Jurkat cells (blue line), permeabilized with 0.5% Triton. Primary incubation overnight (1:50 dilution) followed by Alexa Fluor 488 secondary antibody (1 ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



Immunofluorescence: LHR Antibody [NBP2-76804] - Analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (1:50 dilution) followed by Alexa Fluor 488 secondary antibody (2 ug/ml), showing plasma membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 ug/ml) followed by Alexa Fluor 488 secondary antibody (2 ug/ml).





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-76804

HAF017 Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

291-G1-200 IGF-I/IGF-1 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-76804

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



