# **Product Datasheet**

# Common gamma Chain/IL-2 R gamma Antibody (633162) [Allophycocyanin/Cy7] FAB2842APCCY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/FAB2842APCCY7

Updated 8/3/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/FAB2842APCCY7



# FAB2842APCCY7

Common gamma Chain/IL-2 R gamma Antibody (633162) [Allophycocyanin/Cy7]	
Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark. Do not freeze.
Clonality	Monoclonal
Clone	633162
Preservative	0.05% Sodium Azide
Isotype	lgG2b
Conjugate	Allophycocyanin/Cy7
Purity	Protein A or G purified from hybridoma culture supernatant
Buffer	PBS
Product Description	
Host	Mouse
Gene ID	3561
Gene Symbol	IL2RG
Species	Human
Specificity/Sensitivity	Detects human Common $\gamma$ Chain/IL-2 R $\gamma$ in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human (rh) IL-2 R $\alpha$ , rhIL-2 R $\beta$ , rhIL-15 R $\alpha$ , recombinant mouse IL-2 R $\gamma$ , or recombinant canine IL-2 R $\alpha$ is observed. In Western blots, approximately 80% cross-reactivity with recombinant human IL-2 R $\beta$ , 50% cross-reactivity with rhIL-15 R $\alpha$ and recombinant mouse IL-2 R $\alpha$ , and no cross-reactivity with rhIL-2 R $\alpha$ is observed.
Immunogen	S. frugiperda insect ovarian cell line Sf21-derived recombinant human Common γ Chain/IL-2γ Leu23-Asn254 Accession # P31785
Product Application Details	
Applications	Flow Cytometry
Recommended Dilutions	Flow Cytometry
Application Notes	Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any



questions.



### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

# **Products Related to FAB2842APCCY7**

6507-IL-010/CF IL-4 [Unconjugated]

384-RG-050 Common gamma Chain/IL-2 R gamma [Unconjugated]

202-IL-010 IL-2 [Unconjugated] 247-ILB-005 IL-15 [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/FAB2842APCCY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

