Product Datasheet

PLAG1 Antibody (3B7) - Azide and BSA Free H00005324-M02-50ug

Unit Size: 50 ug

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 20

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/H00005324-M02

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/H00005324-M02



H00005324-M02-50ug

PLAG1 Antibody (3B7) - Azide and BSA Free


50 ug
Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Monoclonal
3B7
No Preservative
IgG2a Kappa
IgG purified
In 1x PBS, pH 7.4

Product Description		
Description	Quality control test: Antibody Reactive Against Recombinant Protein.	
Host	Mouse	
Gene ID	5324	
Gene Symbol	PLAG1	
Species	Human, Mouse	
Immunogen	PLAG1 (NP_002646, 2 a.a. ~ 99 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa. ATVIPGDLSEVRDTQKVPSGKRKRGETKPRKNFPCQLCDKAFNSVEKLKVHSY SHTGERPYKCIQQDCTKAFVSKYKLQRHMATHSPEKTHKCNYCEK	
Notes	This product is produced by and distributed for Abnova, a company based in Taiwan.	

Product Application Details	
Applications	Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:500, ELISA, Immunohistochemistry 1:10-1:500, Immunohistochemistry-Paraffin
Application Notes	Antibody reactivity against recombinant protein with GST tag on ELISA and WB. GST tag alone is used as a negative control. IHC usage was reported in scientific literature.


Images

Western Blot detection against Immunogen (36.52 KDa) .







Left, IHC showing TGFBR3 staining in whole tumor sections (upper panels, scale bars=5mm) and enlargement of selected areas (lower panels, scale bars=100um). Right, quantification of IHC TGFBR3 levels in TGFBR3-PLAG1 positive vs. negative tumors. P (high TGFBR3)=0.002, Fisher's exact test. n=5+34. Image collected and cropped by CiteAb from the following publication (nature.com/articles/s41467-017-01178-z), licensed under a CC-BY license.



Detection limit for recombinant GST tagged PLAG1 is approximately 0.03ng/ml as a capture antibody.



Immunohistochemistry: PLAG1 Antibody (3B7) [H00005324-M02] -Detection of the FGFR1-PLAG1 & TGFBR3-PLAG1 fusion genes. a Illustration of the FGFR1-PLAG1 fusion gene. Arrows show locations of genomic break points for each of the tumors. b Expression of PLAG1 (left) & FGFR1 (right) in PLAG1 fusion-negative vs. FGFR1-PLAG1positive tumors. Graphs show fragments per kilobase of exon per million fragments mapped (FPKM), based on RNA-seq data. ****P < 0.0001; Student's t-test. n = 19 + 7. Horizontal lines show mean values. c Illustration of the TGFBR3-PLAG1 fusion gene. d Expression (FPKM) of PLAG1 (left) & TGFBR3 (right) in PLAG1 fusion-negative vs. TGFBR3-PLAG1-positive tumors. ***P < 0.001, ****P < 0.0001; Student's t-test. n = 19 + 5. Horizontal lines show mean values. e Left, IHC showing TGFBR3 staining in whole tumor sections (upper panels, scale bars=5 mm) & enlargement of selected areas (lower panels, scale bars=100 µm). Right, quantification of IHC TGFBR3 levels in TGFBR3-PLAG1 positive vs. negative tumors. P (high TGFBR3) = 0.002, Fisher's exact test. n = 5 + 34. f Poisson sample clustering based on gene expression in cohort 1, annotated by TGFBR3 IHC results Image collected & cropped by CiteAb from the following publication (https://www.nature.com/articles/s41467-017-01178-z), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Jiarui Li, Yilan Li, Denghui Wang, Rui Liao, Zhongjun Wu PLAG1 interacts with GPX4 to conquer vulnerability to sorafenib induced ferroptosis through a PVT1/miR-195-5p axis-dependent manner in hepatocellular carcinoma Journal of Experimental & Clinical Cancer Research: CR 2024-05-14 [PMID: 38745179]

Sanchez-Avila M, Tjendra Y, Zuo Y et al. Don't SUMP it! Utility of PLAG1 immunocytochemistry in basaloid SUMP subcategory Cancer cytopathology 2023-09-13 [PMID: 37702124] (ICC/IF, Human)

Dalin MG, Katabi N, Persson M et al. Multi-dimensional genomic analysis of myoepithelial carcinoma identifies prevalent oncogenic gene fusions Nature Communications 2017-10-30 [PMID: 29084941]

Anderson WJ, Mariño-Enríquez A, Trpkov K et al. Expanding the Clinicopathologic and Molecular Spectrum of Lipoblastoma-like Tumor in a Series of 28 Cases Modern pathology: an official journal of the United States and Canadian Academy of Pathology, Inc 2023-06-22 [PMID: 37355153]

Adepitan O, Olufunlola A, Oluwole O et al. HMGA2 Immunoexpression is frequent in salivary gland pleomorphic adenoma: immunohistochemical and molecular analyses of PLAG1 and HMGA2 in 25 cases. Int J Clin Exp Pathol. 2022-02-15 [PMID: 35265254]

Sakai H, Fujii Y, Kuwayama N et al. Plag1 regulates neuronal gene expression and neuronal differentiation of neocortical neural progenitor cells. Genes Cells. 2019-10-15 [PMID: 31442350]

Giovacchini F, Bensi C, Belli S et al. Low-grade intraductal carcinoma of salivary glands: A systematic review of this rare entity. J Oral Biol Craniofac Res. 2018-11-25 [PMID: 30555776]

Goto Y, Ibi M, Sato H et al. PLAG1 enhances the stemness profiles of acinar cells in normal human salivary glands in a cell type-specific manner. J Oral Biosci. 2020-01-30 [PMID: 32007659]

Keiko S, Shintaro S, Tomoyuki A et al. Myoepithelioma of soft tissue and bone, and myoepithelioma-like tumors of the vulvar region: Clinicopathological study of 15 cases by PLAG1 immunohistochemistry. Pathol Int. 2020-09-17 [PMID: 32940946]

Yoon YJ, Kim D, Tak KY et al. Salivary gland organoid culture maintains distinct glandular properties of murine and human major salivary glands Nature communications 2022-06-07 [PMID: 35672412] (IF/IHC, Human)

Brcic I, Igrec J, Halbwedl I Et al. Expanding the spectrum of PLAG1-rearranged lipoblastomas arising in patients over 45, with identification of novel fusion partners Modern pathology: an official journal of the United States and Canadian Academy of Pathology, Inc 2021-08-16 [PMID: 34400796]

Gerhard-Hartmann E, Vokuhl C, Roth S Et al. The histological and molecular spectrum of lipoblastoma: A case series with identification of three novel gene fusions by targeted RNA-sequencing Pathology, research and practice 2021-08 -18 [PMID: 34455363] (IHC-P, Human)

More publications at http://www.novusbio.com/H00005324-M02





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to H00005324-M02-50ug

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-96981-0.5mg Mouse IgG2a Kappa Isotype Control (M2AK)

NBP2-56656PEP PLAG1 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/H00005324-M02

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

