Product Datasheet VEGFR1/Flt-1 Antibody (EWC) [HRP] NB600-1004H

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB600-1004H

Updated 7/11/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB600-1004H



NB600-1004H

VEGFR1/Flt-1 Antibody (EWC) [HRP]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clonality Monoclonal Clone EWC Preservative No Preservative Isotype IgG1 Conjugate HRP Purity Protein G purified Buffer PBS Product Description 2321 Gene ID 2321 Gene Symbol FLT1 Species Human, Mouse Reactivity Notes Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB. And MP-2400-NB. Please contact Technical Support if you have any questions. Marker Endothelial Cell Marker Specificity/Sensitivity The monoclonal will detect native and denaturated human VEGFR-1/ FIt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse FIt-1. Immunogen Recombinant human soluble extracellular FIt-1 (D5)	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneEWCPreservativeNo PreservativeIsotypeIgG1ConjugateHRPPurityProtein G purifiedBufferPBSProduct Description2321HostMouseGene ID2321Gene SymbolFLT1SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Unit Size	0.1 ml
ClonalityMonoclonalCloneEWCPreservativeNo PreservativeIsotypeIgG1ConjugateHRPPurityProtein G purifiedBufferPBSProduct Description1000000000000000000000000000000000000	Concentration	•
CloneEWCPreservativeNo PreservativeIsotypeIgG1ConjugateHRPPurityProtein G purifiedBufferPBSProduct DescriptionMouseGene ID2321Gene SymbolFLT1SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Storage	Store at 4C in the dark.
PreservativeNo PreservativeIsotypeIgG1ConjugateHRPPurityProtein G purifiedBufferPBSProduct DescriptionHostMouseGene ID2321Gene SymbolFLT1SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Clonality	Monoclonal
IsotypeIgG1ConjugateHRPPurityProtein G purifiedBufferPBSProduct DescriptionMouseHostMouseGene ID2321Gene SymbolFLT1SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ FIt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse FIt-1.ImmunogenRecombinant human soluble extracellular FIt-1 (D5)	Clone	EWC
ConjugateHRPPurityProtein G purifiedBufferPBSProduct DescriptionMouseHostMouseGene ID2321Gene SymbolFLT1SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ FIt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse FIt-1.ImmunogenRecombinant human soluble extracellular FIt-1 (D5)	Preservative	No Preservative
PurityProtein G purifiedPurityProtein G purifiedBufferPBSProduct DescriptionMouseHostMouseGene ID2321Gene SymbolFLT1SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/FIt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse FIt-1.ImmunogenRecombinant human soluble extracellular FIt-1 (D5)	Isotype	IgG1
Buffer PBS Product Description Host Mouse Gene ID 2321 Gene Symbol FLT1 Species Human, Mouse Reactivity Notes Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions. Marker Endothelial Cell Marker Specificity/Sensitivity The monoclonal will detect native and denaturated human VEGFR-1/ FIt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse FIt-1. Immunogen Recombinant human soluble extracellular FIt-1 (D5)	Conjugate	HRP
Product Description Host Mouse Gene ID 2321 Gene Symbol FLT1 Species Human, Mouse Reactivity Notes Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions. Marker Endothelial Cell Marker Specificity/Sensitivity The monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1. Immunogen Recombinant human soluble extracellular Flt-1 (D5)	Purity	Protein G purified
HostMouseGene ID2321Gene SymbolFLT1SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Buffer	PBS
Gene ID2321Gene SymbolFLT1SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Product Description	
Gene Symbol FLT1 Species Human, Mouse Reactivity Notes Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions. Marker Endothelial Cell Marker Specificity/Sensitivity The monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1. Immunogen Recombinant human soluble extracellular Flt-1 (D5)	Host	Mouse
SpeciesHuman, MouseReactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Gene ID	2321
Reactivity NotesPlease note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Gene Symbol	FLT1
host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.MarkerEndothelial Cell MarkerSpecificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Species	Human, Mouse
Specificity/SensitivityThe monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1.ImmunogenRecombinant human soluble extracellular Flt-1 (D5)	Reactivity Notes	host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact
ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1. Immunogen Recombinant human soluble extracellular Flt-1 (D5)	Marker	Endothelial Cell Marker
	Specificity/Sensitivity	ELISA experiments and on the surface of different human cell types. The will
Product Application Details	Immunogen	Recombinant human soluble extracellular Flt-1 (D5)
· · · · · · · · · · · · · · · · · · ·	Product Application Details	
Applications Western Blot, ELISA, Immunocytochemistry/Immunofluorescence, In vivo assay	Applications	Western Blot, ELISA, Immunocytochemistry/Immunofluorescence, In vivo assay
Recommended Dilutions Western Blot, ELISA, Immunocytochemistry/Immunofluorescence, In vivo assay	Recommended Dilutions	Western Blot, ELISA, Immunocytochemistry/Immunofluorescence, In vivo assay
Application NotesOptimal dilution of this antibody should be experimentally determined.	Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NB600-1004H

NBP1-97074	Mouse IgG1 Isotype Control (MG1) [HRP]
NB100-527PEP	VEGFR1/FIt-1 Antibody Blocking Peptide
DVE00	VEGF [HRP]
DVR100C	VEGFR1/Flt-1 [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-1004H

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

