Product Datasheet

Goat anti-Hamster IgG (H+L) Secondary Antibody [PE/Cy5.5] (Pre-adsorbed) NBP1-71730PECY55

Unit Size: 500 ul

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-71730PECY55

Updated 7/11/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP1-71730PECY55



NBP1-71730PECY55

Goat anti-Hamster IgG (H+L) Secondary Antibody [PE/Cy5.5] (Pre-adsorbed)

Goat anti-Hamster IgG (H+L) Secondary Antibody [PE/Cy5.5] (Pre-adsorbed)	
Product Information	
Unit Size	500 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark. Do not freeze.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Conjugate	PE/Cy5.5
Purity	Multi-step
Buffer	PBS
Product Description	
Host	Goat
Species	Hamster
Reactivity Notes	Cricetulus migratorius (Armenian Hamster); Melanochromis auratus (Golden Syrian Hamster)
Specificity/Sensitivity	This antibody was pre-adsorbed against Mouse and Rat Serum Proteins. No reaction was observed against Mouse or Rat serum proteins. Specificity was confirmed using ELISA at less than 1% cross reactivity.
Immunogen	Armenian and Golden Syrian Hamster IgG, whole molecule
Product Application Details	
Applications	Western Blot, Dot Blot, ELISA, Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot, ELISA, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Dot Blot, Fluorophore-linked immunosorbent assay
Application Notes	Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any



questions.



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-71730PECY55

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

