# **Product Datasheet**

# Bcl-2 Antibody (SPM117) [CoraFluor™ 1] NBP2-34440CL1

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34440CL1

Updated 8/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34440CL1



## NBP2-34440CL1

Bcl-2 Antibody (SPM117) [CoraFluor™ 1]

Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1, thiol reactive  For more information, please see our CoraFluor(TM) TR-FRET technology flyer Mouse  Gene ID  596  Gene Symbol  BCL2  Species  Human, Mouse (Negative), Rat (Negative)  Peactivity Notes  Does not react with Mouse or Rat.  Marker  Apoptosis & Follicular Lymphoma Marker  Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers are negative Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.	BCI-2 Antibody (SPIVITIT) [C	orafiuor ···· Ij	
Please see the vial label for concentration. If unlisted please contact technical services.	Product Information		
Storage Store at 4C in the dark. Do not freeze.  Clonality Monoclonal  Clone SPM117  Preservative No Preservative  Isotype IgG1 Kappa  Conjugate CoraFluor 1  Purity Protein A or G purified  Buffer PBS  Product Description  Description CaraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, s45 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 and be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1. amine reactive  CoraFluor(TM) 1. amine reactive  CoraFluor(TM) 1. amine reactive  CoraFluor(TM) 1. The reactive  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death for centers are negative and neoplastic enters are negative and neoplastic cell sair beneative and neoplastic cell sair centers are negative and neoplastic cell	Unit Size	0.1 ml	
Clone SPM117  Preservative No Preservative  Igot Kappa  Conjugate CoraFluor 1  Purity Protein A or G purified  Buffer PBS  Product Description  CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolve Fluorescence Resonance Energy Transfer) or TRF (Time-Resolve Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1, thiol reactive  For more information, please see our CoraFluor(TM) TR-FRET technology flyer  Host Mouse  Gene ID 596  Gene Symbol BCL2  Species Human, Mouse (Negative), Rat (Negative)  Reactivity Notes Does not react with Mouse or Rat.  Marker Apoptosis & Follicular Lymphoma Marker  Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular rymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 regetive.	Concentration	·	
Clone SPM117  Preservative No Preservative  Isotype IgG1 Kappa Conjugate CoraFluor 1  Purity Protein A or G purified  Buffer PBS  Product Description  CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolve Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, temary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1, thiol reactive  For more information, please see our CoraFluor(TM) TR-FRET technology flyer Mouse  Gene ID 596  Gene Symbol BCL2  Species Human, Mouse (Negative), Rat (Negative)  Reactivity Notes Does not react with Mouse or Rat.  Marker Apoptosis & Follicular Lymphoma Marker  Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein, whereas the normal or hyperplastic germinal centers are negative Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular religifieration in lymph node biopsies. It may also be use in distinguishing between reactive and the small number in which the neoplastic cells are Bcl-2 regative.  Immunogen	Storage	Store at 4C in the dark. Do not freeze.	
Preservative   Isotype   IgG1 Kappa	Clonality	Monoclonal	
IgG1 Kappa   IgG1 Kappa   Conjugate   CoraFluor 1	Clone	SPM117	
Conjugate  CoraFluor 1  Purity  Protein A or G purified  Buffer  PBS  Product Description  CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolve Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1. thiol reactive  For more information, please see our CoraFluor(TM) TR-FRET technology flyer  Host Mouse  Gene ID 596  Gene Symbol BCL2  Species Human, Mouse (Negative), Rat (Negative)  Reactivity Notes Does not react with Mouse or Rat.  Marker Apoptosis & Follicular Lymphoma Marker  Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this artibody is valuable when distinguishing between reactive and neoplastic follicular profileration in lymph noab petween exactive and neoplastic follicular profileration in lymph noab petween reactive and neoplastic follicular profileration in lymph noab petween reactive and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen	Preservative	No Preservative	
Purity Protein A or G purified  Buffer PBS  Product Description  CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1, thiol reactive  For more information, please see our CoraFluor(TM) TR-FRET technology flyer Mouse  Gene ID  596  Gene Symbol  BCL2  Species  Human, Mouse (Negative), Rat (Negative)  Reactivity Notes  Does not react with Mouse or Rat.  Marker  Apoptosis & Follicular Lymphoma Marker  Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular profileration in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen	Isotype	IgG1 Kappa	
PBS	Conjugate	CoraFluor 1	
Product Description  CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 coras be used for the development of robust and scalab TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1. amine reactive  CoraFluor(TM) 1. thiol reactive  For more information, please see our CoraFluor(TM) TR-FRET technology flyer Mouse  Gene ID  596  Gene Symbol  BCL2  Species  Human, Mouse (Negative), Rat (Negative)  Reactivity Notes  Does not react with Mouse or Rat.  Apoptosis & Follicular Lymphoma Marker  Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen	Purity	Protein A or G purified	
CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.    CoraFluor(TM) 1, amine reactive	Buffer	PBS	
Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1, amine reactive  CoraFluor(TM) 1, thiol reactive  For more information, please see our CoraFluor(TM) TR-FRET technology flyer Mouse  Gene ID  596  Gene Symbol  BCL2  Species  Human, Mouse (Negative), Rat (Negative)  Reactivity Notes  Does not react with Mouse or Rat.  Marker  Apoptosis & Follicular Lymphoma Marker  Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-v or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 regative.  Immunogen	Product Description		
Host Gene ID 596 Gene Symbol BCL2 Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Apoptosis & Follicular Lymphoma Marker Specificity/Sensitivity This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers are negative Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen  A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein	Description	Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.  CoraFluor(TM) 1, amine reactive	
Gene SymbolBCL2SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerApoptosis & Follicular Lymphoma MarkerSpecificity/SensitivityThis antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.ImmunogenA synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein	Host		
Species  Human, Mouse (Negative), Rat (Negative)  Reactivity Notes  Does not react with Mouse or Rat.  Marker  Apoptosis & Follicular Lymphoma Marker  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen  A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein	Gene ID	596	
Reactivity Notes  Does not react with Mouse or Rat.  Marker  Apoptosis & Follicular Lymphoma Marker  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative.  Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen  A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein	Gene Symbol	BCL2	
Marker  Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen  A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein	Species	Human, Mouse (Negative), Rat (Negative)	
Specificity/Sensitivity  This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen  A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein	Reactivity Notes	Does not react with Mouse or Rat.	
oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be use in distinguishing between those follicular lymphomas that express Bcl-2 protein and the small number in which the neoplastic cells are Bcl-2 negative.  Immunogen  A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein	Marker	Apoptosis & Follicular Lymphoma Marker	
	Specificity/Sensitivity	oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of Bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of Bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be used in distinguishing between those follicular lymphomas that express Bcl-2 protein	
	Immunogen	A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein. (Uniprot: P10415)	



Notes	CoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, CyTOF-ready
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP2-34440CL1**

NBP2-34889-100ug Recombinant Human Bcl-2 Protein

210-TA-005 TNF-alpha [Unconjugated]

DYC827B-2 Bcl-2 [Biotin]

AF835 Caspase-3 Antibody [Unconjugated] - Active

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34440CL1

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

