

Product Datasheet

Thyroglobulin Antibody (2H11 + 6E1) [Janelia Fluor® 585] NBP2-34530JF585

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-34530JF585

Updated 8/15/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-34530JF585



NBP2-34530JF585

Thyroglobulin Antibody (2H11 + 6E1) [Janelia Fluor® 585]

| Product Information | |
|---------------------|---|
| Unit Size | 0.1 ml |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at 4C in the dark. |
| Clonality | Monoclonal |
| Clone | 2H11 + 6E1 |
| Preservative | 0.05% Sodium Azide |
| Isotype | IgG1 Kappa/IgG1 Kappa |
| Conjugate | Janelia Fluor 585 |
| Purity | Protein A or G purified |
| Buffer | 50mM Sodium Borate |

| Product Description | |
|-------------------------|---|
| Host | Mouse |
| Gene ID | 7038 |
| Gene Symbol | TG |
| Species | Human, Mouse, Rat |
| Marker | Thyroidal Cell Marker |
| Specificity/Sensitivity | Thyroglobulin is a 660kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo conformational maturation in the endoplasmic reticulum. The vast majority of follicular carcinomas of the thyroid will give positive immunoreactivity for anti-thyroglobulin even though sometimes only focally. Poorly differentiated carcinomas of the thyroid are frequently anti-thyroglobulin negative. Adenocarcinomas of other-than-thyroid origin do not react with this antibody. This antibody is useful in identification of thyroid carcinoma of the papillary and follicular types. Presence of thyroglobulin in metastatic lesions establishes the thyroid origin of tumor. Anti-thyroglobulin, combined with anti-calcitonin, can identify medullary carcinomas of the thyroid. Furthermore, anti-thyroglobulin, combined with anti-TTF1, can be a reliable marker to differentiate between primary thyroid and lung neoplasms. |
| Immunogen | Human thyroid follicular cells |
| Notes | Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. |

| Product Application Details | |
|-----------------------------|--|
| Applications | Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready |
| Recommended Dilutions | Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready |
| Application Notes | Optimal dilution of this antibody should be experimentally determined. |





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-
techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis.
Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34530JF585

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

