Product Datasheet

Cytokeratin 8/18 Antibody (K8.8 + DC10) - Azide and BSA Free NBP2-34655-0.1mg

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34655

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-34655



NBP2-34655-0.1mg


Cytokeratin 8/18 Antibody (K8.8 + DC10) - Azide and BSA Free

| Cytokeratin 8/18 Antibody (K8.8 + DC10) - Azide and BSA Free | |
|--|--|
| Product Information | |
| Unit Size | 0.1 mg |
| Concentration | 1.0 mg/ml |
| Storage | Store at -20 to -80C. Avoid freeze-thaw cycles. |
| Clonality | Monoclonal |
| Clone | K8.8 + DC10 |
| Preservative | No Preservative |
| Isotype | IgG1 Kappa/IgG1 Kappa |
| Purity | Protein A or G purified |
| Buffer | 10 mM PBS |
| Product Description | |
| Description | 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-34344). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. |
| Host | Mouse |
| Gene ID | 3856 |
| Gene Symbol | KRT8 |
| Species | Human |
| Specificity/Sensitivity | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody cocktail recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the normal epidermis so only Paget cells are stained. Immunohistochemical staining with this monoclonal antibody is indistinguishable from that obtained with monoclonal antibody 5D3. |
| Immunogen | Keratin preparation from a human carcinoma (K8.8); PMC-42 human breast carcinoma cells (DC10) |
| Product Application Details | |
| Applications | Western Blot, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, CyTOF-ready |
| Recommended Dilutions | Western Blot 0.5-1.0ug/ml, Flow Cytometry 0.5-1ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2ug/ml, Immunohistochemistry-Paraffin 0.5-1.0ug/ml, Immunohistochemistry-Frozen 0.5-1.0ug/ml, Flow (Intracellular), CyTOF-ready |
| Application Notes | Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined. |




Images

Immunohistochemistry-Paraffin: Cytokeratin 8/18 Antibody (K8.8 + DC10) - Azide and BSA Free [NBP2-34655] - Formalin-fixed paraffinembedded human colon carcinoma stained with cytokeratin 8/18 Monoclonal Antibody (K8.8 +DC10)



Immunohistochemistry-Paraffin: Cytokeratin 8/18 Antibody (K8.8 + DC10) - Azide and BSA Free [NBP2-34655] - Formalin-fixed, paraffinembedded human Overian cancer stained with Cytokeratin 8/18 Monoclonal Antibody (K8.8 + DC10).



Publications

Shenderov E, De Marzo AM, Lotan TL et al. Neoadjuvant enoblituzumab in localized prostate cancer: a single-arm, phase 2 trial Nature medicine 2023-04-01 [PMID: 37012549]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34655-0.1mg

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

8184-CK-050 Choline Kinase beta [Unconjugated]

MAB1368 alpha-Fetoprotein/AFP Antibody (189502) [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34655

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

