Product Datasheet

Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody (M2-7C10 + M2-9E3 + T311 + HMB45) - Azide and BSA Free NBP2-34681-0.1mg

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34681

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34681



NBP2-34681-0.1mg

Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody (M2-7C10 + M2-9E3 + T311 + HMB45) - Azide and BSA Free		
Product Information		
Unit Size	0.1 mg	
Concentration	1.0 mg/ml	
Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.	
Clonality	Monoclonal	
Clone	M2-7C10 + M2-9E3 + T311 + HMB45	
Preservative	No Preservative	
Isotype	IgG2b Lambda/IgG2a Kappa/IgG1 Kappa	
Purity	Protein A or G purified	
Buffer	10 mM PBS	
Product Description		
Description	1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-34337). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.	
Host	Mouse	
Gene ID	2315	
Gene Symbol	MLANA	
Species	Human	

Gene Symbol	MLANA
Species	Human

Marker	Melanoma Marker
iidi itoi	molarionia markoi

Specificity/Sensitivity

This antibody cocktail recognizes three melanoma-specific proteins, which include MART-1, Tyrosinase and gp100. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Tyrosinase is one of the targets for cytotoxic T-cell recognition in melanoma patients. Function of gp100 is not known but it is reported to be a useful marker for melanocytes and melanomas. This cocktail of three markers is designed for extremely sensitive labeling of formalin-fixed, paraffin-embedded melanomas and other tumors showing melanocytic differentiation.


	9
Immunogen	Recombinant Melanoma Marker (MART-1 + Tyrosinase + gp100) protein (M2-7C10; M2-9E3); Recombinant tyrosinase protein (T311); Extract of pigmented melanoma metastases from lymph nodes (HMB45)

	meianoma metastases from lymph nodes (HIMB45)
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 2-4 ug/ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 0.5-1.0ug/ml, Immunohistochemistry-Frozen 0.5-1.0ug/ml
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.




Images


Western Blot: Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody (M2-7C10 + M2-9E3 + T311 + HMB45) - Azide and BSA Free [NBP2-34681] - Western Blot Analysis of COLO38 cell lysate using Melanoma Marker (MART-1 + Tyrosinase + gp100) MAb (M2-7C10 + M2 -9E3 + T311 + HMB45).




Immunohistochemistry-Frozen: Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody (M2-7C10 + M2-9E3 + T311 + HMB45) - Azide and BSA Free [NBP2-34681] - Frozen human melanoma stained for melanoma cancer cells (red) and counter-stained with DAPI (blue). This image was submitted via customer review. Image from the Alexa Fluor 700 version of this antibody.




Immunohistochemistry-Frozen: Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody (M2-7C10 + M2-9E3 + T311 + HMB45) - Azide and BSA Free [NBP2-34681] - Melanoma Tissue stained for Mart-1 Tyrosinase p100 (red) counterstained with DAPI (blue). Image from a verified customer review.



Immunohistochemistry-Paraffin: Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody (M2-7C10 + M2-9E3 + T311 + HMB45) - Azide and BSA Free [NBP2-34681] - Formalin-fixed, paraffin-embedded human melanoma stained with Melanoma Marker MAb (M2-7C10 + M2-9E3 + T311 + HMB45).



Immunohistochemistry: Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody (M2-7C10 + M2-9E3 + T311 + HMB45) - Azide and BSA Free [NBP2-34681] - Scan of human melanoma sample. Stained for Mart-1+Tyrosinase+gp100 (green) counter stained with DAPI (blue). Image from verified customer review. Image using the Janelia Fluor 549 format of this antibody.



Publications

Liu Q, Adhikari E, Lester D et al. Androgen Drives Melanoma Invasiveness and Metastatic Capacity by Promoting Tumorigenic Fucosylation Research Square 2023-03-10 (IHC-P)

Martinek J, Lin J, Kim KI Et al. Transcriptional profiling of macrophages in situ in metastatic melanoma reveals localization-dependent phenotypes and function Cell Rep Med 2022-05-18 [PMID: 35584631]

Details:

Citation using the Azide and BSA Free version of this antibody.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-34681-0.1mg

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

285-IF-100 IFN-gamma [Unconjugated]

202-IL-010 IL-2 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34681

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

