

# Product Datasheet

## MUC1 Antibody (SPM533) [Alexa Fluor™ Plus 680] NBP2-34782AFP680

Unit Size: 0.1 ml

Store at 4C in the dark.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-34782AFP680](http://www.novusbio.com/NBP2-34782AFP680)

Updated 4/12/2026 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-34782AFP680](http://www.novusbio.com/reviews/destination/NBP2-34782AFP680)



**NBP2-34782AFP680**

MUC1 Antibody (SPM533) [Alexa Fluor™ Plus 680]

| Product Information |   |
|---------------------|---|
| Unit Size           | 0.1 ml  |
| Concentration       | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage             | Store at 4C in the dark.  |
| Clonality           | Monoclonal  |
| Clone               | SPM533  |
| Preservative        | 0.05% Sodium Azide  |
| Isotype             | IgG2a Lambda  |
| Conjugate           | Alexa Fluor Plus 680  |
| Purity              | Protein A or G purified   |
| Buffer              | 50mM Sodium Borate  |

| Product Description     |  |
|-------------------------|--|
| Host                    | Mouse  |
| Gene ID                 | 4582   |
| Gene Symbol             | MUC1   |
| Species                 | Human  |
| Marker                  | Epithelial Marker  |
| Specificity/Sensitivity | In Western blotting, it recognizes proteins in MW range of 265-400kDa, identified as different glycoforms of EMA. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit and the C-terminal beta subunit. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. EMA may provide a protective layer on epithelial cells against bacterial and enzyme attack. The beta subunit contains a C-terminal domain, which is involved in cell signaling, through phosphorylations and protein-protein interactions. In immunohistochemical assays, it superbly stains routine formalin/paraffin carcinomas. Antibody to EMA is useful as a pan-epithelial marker for detecting early metastatic loci of carcinoma in bone marrow or liver.   |
| Immunogen               | Delipidated extract of human milk fat globule membranes  |
| Notes                   | This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its components for resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or <a href="mailto:outlicensing@thermofisher.com">outlicensing@thermofisher.com</a> . This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. |

**Product Application Details**

|                              |  |
|------------------------------|--|
| <b>Applications</b>          | Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/<br>Immunofluorescence, Immunohistochemistry, CyTOF-ready |
| <b>Recommended Dilutions</b> | Flow Cytometry, Immunohistochemistry, Immunocytochemistry/<br>Immunofluorescence, Immunohistochemistry-Paraffin, CyTOF-ready |
| <b>Application Notes</b>     | Optimal dilution of this antibody should be experimentally determined.   |





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP2-34782AFP680**

---

|                    |   |
|--------------------|---|
| H00004582-Q01-10ug | Recombinant Human MUC1 GST (N-Term) Protein |
| 1129-ER-050        | ErbB2/Her2 [Unconjugated]                   |
| 10332-MU-050       | MUC1 [Unconjugated]                         |
| NB300-141          | GFAP Antibody                               |

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-34782AFP680](http://www.novusbio.com/reviews/submit/NBP2-34782AFP680)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

