## **Product Datasheet**

# CD45RA Antibody (SPM568) - Azide and BSA Free NBP2-34802-0.1mg

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-34802

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-34802



## NBP2-34802-0.1mg


CD45RA Antibody (SPM568) - Azide and BSA Free

| CD45RA Antibody (SPM568) - Azide and BSA Free |   |
|---|---|
| Product Information                           |   |
| Unit Size                                     | 0.1 mg  |
| Concentration                                 | 1.0 mg/ml   |
| Storage                                       | Store at -20 to -80C. Avoid freeze-thaw cycles.   |
| Clonality                                     | Monoclonal  |
| Clone   | SPM568  |
| Preservative                                  | No Preservative   |
| Isotype                                       | IgG2a Kappa   |
| Purity  | Protein A or G purified   |
| Buffer  | 10 mM PBS   |
| Product Description                           |   |
| Description                                   | 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-32936).  Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.   |
| Host  | Mouse   |
| Gene ID                                       | 5788  |
| Gene Symbol                                   | PTPRC   |
| Species                                       | Human   |
| Marker  | Leucocyte Marker  |
| Specificity/Sensitivity                       | Recognizes a protein of 205kDa-220kDa, identified as CD45RA. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve). |
| Immunogen                                     | Stimulated human leukocytes   |
| Product Application Details                   |   |
| Applications                                  | Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin  |
| Recommended Dilutions                         | Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin   |
| Application Notes                             | Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.   |




#### **Images**


Western Blot: CD45RA Antibody (SPM568) - Azide and BSA Free [NBP2 -34802] - Western Blot Analysis of human spleen tissue lysates using CD45RA Antibody (SPM568).




Immunocytochemistry/Immunofluorescence: CD45RA Antibody (SPM568) - Azide and BSA Free [NBP2-34802] - Immunofluorescence Analysis of PFA-fixed Jurkat cells labeling CD45RA with CD45RA Antibody (SPM568) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Red Dot.



Immunohistochemistry-Paraffin: CD45RA Antibody (SPM568) - Azide and BSA Free [NBP2-34802] - Formalin-paraffin human spleen stained with CD45RA MAb (SPM568).



Flow Cytometry: CD45RA Antibody (SPM568) - Azide and BSA Free [NBP2-34802] - Flow Cytometric Analysis of Jurkat cells using CD45RA Antibody (SPM568) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).





#### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP2-34802-0.1mg

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-96981-0.5mg Mouse IgG2a Kappa Isotype Control (M2AK)

NBP2-34802AF405 CD45RA Antibody (SPM568) [Alexa Fluor® 405]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-34802

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

