Product Datasheet

AMPD3 Antibody (AMPD3/901) [Janelia Fluor® 525] NBP2-47691JF525

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47691JF525

Updated 11/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-47691JF525



NBP2-47691JF525

AMPD3 Antibody (AMPD3/901) [Janelia Fluor® 525]

lia Fluor® 525]
ml
ase see the vial label for concentration. If unlisted please contact technical ices.
e at 4C in the dark.
oclonal
PD3/901
% Sodium Azide
2b Kappa
elia Fluor 525
ein A or G purified
M Sodium Borate
ise
PD3
nan
hroid Marker
cognizes a protein of ~90kDa, which is identified as Adenosine apphosphate Deaminase, isoform E (AMPD3). It has 767 amino acids and is gned an EC 3.5.4.6. It is a highly regulated enzyme that catalyzes the olytic deamination of adenosine monophosphate to inosine monophosphate, anch point in the adenylate catabolic pathway. AMPD3 gene encodes the procyte (E) isoforms, whereas other family members encode isoforms that dominate in muscle (M) and liver (L) cells. This monoclonal antibody shows stivity with cells of the erythroid lineage at all stages of maturation in the otheral blood, bone marrow, and fetal liver. Non-erythroid lineages are ative by flow cytometry. This monoclonal antibody is useful in the diagnosis rythroleukemia, identification of bone marrow erythroid precursors, gating hroid nucleated precursor cells from malignant cells in bone marrow
cognizes a protein of ~90kDa, which is identified as Adenosine hophosphate Deaminase, isoform E (AMPD3). It has 767 amino acids and is gned an EC 3.5.4.6. It is a highly regulated enzyme that catalyzes the rolytic deamination of adenosine monophosphate to inosine monophosphate, anch point in the adenylate catabolic pathway. AMPD3 gene encodes the hrocyte (E) isoforms, whereas other family members encode isoforms that dominate in muscle (M) and liver (L) cells. This monoclonal antibody shows ativity with cells of the erythroid lineage at all stages of maturation in the otheral blood, bone marrow, and fetal liver. Non-erythroid lineages are ative by flow cytometry. This monoclonal antibody is useful in the diagnosis rythroleukemia, identification of bone marrow erythroid precursors, gating hroid nucleated precursor cells from malignant cells in bone marrow cimens.
cognizes a protein of ~90kDa, which is identified as Adenosine iophosphate Deaminase, isoform E (AMPD3). It has 767 amino acids and is gned an EC 3.5.4.6. It is a highly regulated enzyme that catalyzes the rolytic deamination of adenosine monophosphate to inosine monophosphate, anch point in the adenylate catabolic pathway. AMPD3 gene encodes the procyte (E) isoforms, whereas other family members encode isoforms that dominate in muscle (M) and liver (L) cells. This monoclonal antibody shows entivity with cells of the erythroid lineage at all stages of maturation in the otheral blood, bone marrow, and fetal liver. Non-erythroid lineages are active by flow cytometry. This monoclonal antibody is useful in the diagnosis rythroleukemia, identification of bone marrow erythroid precursors, gating hroid nucleated precursor cells from malignant cells in bone marrow cimens. Ombinant full-length human AMPD3 protein (Uniprot: Q01432)
cognizes a protein of ~90kDa, which is identified as Adenosine iophosphate Deaminase, isoform E (AMPD3). It has 767 amino acids and is gned an EC 3.5.4.6. It is a highly regulated enzyme that catalyzes the rolytic deamination of adenosine monophosphate to inosine monophosphate, anch point in the adenylate catabolic pathway. AMPD3 gene encodes the procyte (E) isoforms, whereas other family members encode isoforms that dominate in muscle (M) and liver (L) cells. This monoclonal antibody shows entivity with cells of the erythroid lineage at all stages of maturation in the otheral blood, bone marrow, and fetal liver. Non-erythroid lineages are active by flow cytometry. This monoclonal antibody is useful in the diagnosis rythroleukemia, identification of bone marrow erythroid precursors, gating hroid nucleated precursor cells from malignant cells in bone marrow cimens. Ombinant full-length human AMPD3 protein (Uniprot: Q01432)
cognizes a protein of ~90kDa, which is identified as Adenosine sophosphate Deaminase, isoform E (AMPD3). It has 767 amino acids and is gned an EC 3.5.4.6. It is a highly regulated enzyme that catalyzes the olytic deamination of adenosine monophosphate to inosine monophosphate, anch point in the adenylate catabolic pathway. AMPD3 gene encodes the procyte (E) isoforms, whereas other family members encode isoforms that dominate in muscle (M) and liver (L) cells. This monoclonal antibody shows strivity with cells of the erythroid lineage at all stages of maturation in the oberal blood, bone marrow, and fetal liver. Non-erythroid lineages are active by flow cytometry. This monoclonal antibody is useful in the diagnosis rythroleukemia, identification of bone marrow erythroid precursors, gating hroid nucleated precursor cells from malignant cells in bone marrow cimens. The process of the erythroid process of the diagnosis of the erythroid precursors are gative by flow cytometry. This monoclonal antibody is useful in the diagnosis of the erythroleukemia, identification of bone marrow erythroid precursors, gating throid nucleated precursor cells from malignant cells in bone marrow cimens. The process of the erythroid precursors are gative by flow cytometry. This monoclonal antibody is useful in the diagnosis rythroleukemia, identification of bone marrow erythroid precursors, gating throid nucleated precursor cells from malignant cells in bone marrow cimens.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-47691JF525

NBP1-43317JF525 Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Janelia Fluor

525]

NBP2-14290PEP AMPD3 Recombinant Protein Antigen

210-TA-005 TNF-alpha [Unconjugated]

NBP2-08635 AMPD3 Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47691JF525

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

