# **Product Datasheet**

## Calponin 1 Antibody (CNN1/832 + CALP) [Alexa Fluor® 350] NBP2-47758AF350

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-47758AF350

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-47758AF350

## NBP2-47758AF350

Calponin 1 Antibody (CNN1/832 + CALP) [Alexa Fluor® 350]

Product Information     Unit Size   0.1 ml     Concentration   Please see the vial label for concentration. If unlisted please contact technical services.     Storage   Store at 4C in the dark.     Clonality   Monoclonal     Clone   CNN1/832 + CALP     Preservative   0.05% Sodium Azide     Isotype   IgG1 Kappa     Conjugate   Alexa Fluor 350     Purity   Protein A or G purified     Buffer   50mM Sodium Borate     Product Description   Mouse     Gene ID   1264     Gene ID   1264     Gene Symbol   CNN1     Species   Human, Rat     Marker   Smooth Muscle Marker     Specificity/Sensitivity   Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa of m. Expression of the 29kDa form. I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blothing, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of numan aortic medial smooth muscle. In Western blothing, this monoclonal in vascular and visceral smooth muscle of the uroprovoxin binding protein, which is thought to be in		, <b>.</b> .			
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1264Gene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital artact, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of culturated human foreskin. Calponin is restricted to smooth muscle extra of und visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of unuan aortic antibody reacts with only the 34kDa form of calponin in extracts of unuan aortic in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody moth muscle and is unreactive with fibroblast extracts of unuan aortic antibody moth muscle. Calponin is restricted to smooth muscle contraction. Calponin expression is restricted to smooth muscle of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Product Information				
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1264HostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Unit Size	0.1 ml			
ClonalityMonoclonalCloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1264Gene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34KDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. Im Western blotting, this monoclonal antibody reacts with only the 34kDa form of smooth muscle during which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth cultived human anotic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin in serticted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Concentration				
CloneCNN1/832 + CALPPreservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form, Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and viscerial smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin expression is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Storage	Store at 4C in the dark.			
Preservative 0.05% Sodium Azide   Isotype IgG1 Kappa   Conjugate Alexa Fluor 350   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description Host   Host Mouse   Gene ID 1264   Gene Symbol CNN1   Species Human, Rat   Marker Smooth Muscle Marker   Specificity/Sensitivity Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin is a craamodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.   Immunogen Recombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Clonality	Monoclonal			
IsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and troponyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Clone	CNN1/832 + CALP			
ConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is restricted to smooth muscle contraction. Calponin expression is restricted to smooth muscle contraction. Calponin expression is restricted to smooth muscle contraction. Calponin expression is restricted to smooth muscle contraction. Calponin is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Preservative	0.05% Sodium Azide			
Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description    Host Mouse   Gene ID 1264   Gene Symbol CNN1   Species Human, Rat   Marker Smooth Muscle Marker   Specificity/Sensitivity Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, 1-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa for of calponin in extracts of numan aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.   Immunogen Recombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Isotype	IgG1 Kappa			
Buffer 50mM Sodium Borate   Product Description Host   Host Mouse   Gene ID 1264   Gene Symbol CNN1   Species Human, Rat   Marker Smooth Muscle Marker   Specificity/Sensitivity Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of numan aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.   Immunogen Recombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Conjugate	Alexa Fluor 350			
Product Description   Host Mouse   Gene ID 1264   Gene Symbol CNN1   Species Human, Rat   Marker Smooth Muscle Marker   Specificity/Sensitivity Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of numan aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.   Immunogen Recombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Purity	Protein A or G purified			
HostMouseGene ID1264Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Buffer	50mM Sodium Borate			
Gene ID 1264   Gene Symbol CNN1   Species Human, Rat   Marker Smooth Muscle Marker   Specificity/Sensitivity Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is urreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.   Immunogen Recombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Product Description				
Gene SymbolCNN1SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Host	Mouse			
SpeciesHuman, RatMarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Gene ID	1264			
MarkerSmooth Muscle MarkerSpecificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Gene Symbol	CNN1			
Specificity/SensitivityMultiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Species	Human, Rat			
molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.ImmunogenRecombinant full-length human Calponin 1 protein (Calponin 1/832); Crude	Marker	Smooth Muscle Marker			
	Specificity/Sensitivity	molecular weight isoforms exist; a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, whereas the higher molecular weight variant has been demonstrated in vascular and visceral smooth muscle. In Western blotting, this monoclonal antibody reacts with only the 34kDa form of calponin in extracts of human aortic medial smooth muscle and is unreactive with fibroblast extracts of cultivated human foreskin. Calponin is a calmodulin, F-actin and tropomyosin binding protein, which is thought to be involved in the regulation of smooth muscle contraction. Calponin expression is restricted to smooth muscle cells and has been shown to be a marker of the differentiated (contractile) phenotype of developing smooth muscle.			
	Immunogen				

www.novusbio.com



Notes
-------

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

#### **Product Application Details**

• •	
	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

#### Images

Calponin 1 Antibody (CNN1/832 + CALP) [Alexa Fluor® 350] [NBP2-47758AF350] - Vial of Alexa Fluor 350 conjugated antibody. Alexa Fluor 350 is optimally excited at 346 nm by the UV laser (350 or 355 nm) and has an emission maximum of 442 nm.

	2755) (SA2			
	Alexa Fluc	or® 350		
	LASER (nm)	FILTER	]	
lexa Fluore 350	LASER (nm) UV (350)	FILTER 450/45		
Nexa Fluor® 350	the second se		]	
llexa Fluor <sup>e</sup> 35 <sup>0</sup>	the second se	450/45 EMISSION MAX (nm)	]	
Nexa Fluor® 350	UV (350)	450/45		





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

## **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-47758AF350

Recombinant Human Calponin 1 His Protein
ErbB2/Her2 [Unconjugated]
Calponin 1 Overexpression Lysate
CD31/PECAM-1 Antibody [Unconjugated]
E

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-47758AF350

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

