# **Product Datasheet**

# Complement Factor H Antibody (028B-244.2.10X (10-10)) [mFluor Violet 610 SE] NBP2-52743MFV610

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-52743MFV610

Updated 7/30/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-52743MFV610



#### NBP2-52743MFV610

Complement Factor H Antibody (028B-244.2.10X (10-10)) [mFluor Violet 610 SE]	
Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark.
Clonality	Monoclonal
Clone	028B-244.2.10X (10-10)
Preservative	0.05% Sodium Azide
Isotype	IgG1
Conjugate	mFluor Violet 610 SE
Purity	Protein A purified
Buffer	50mM Sodium Borate
Product Description	
Host	Mouse
Gene ID	3075
Gene Symbol	CFH
Species	Human
Reactivity Notes	Reacts with Human.
Specificity/Sensitivity	NB100-64342 recognizes human complement factor H which exists in 2 forms.
	The most common form, of 150 kDa, and the less common form of 43 kDa. Factor H is secreted by the liver into the blood serum. It is important in regulating the complement pathway, preventing unnecessary inflammation which can damage the host tissue. Complement factor H functions as a cofactor in the inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.
Immunogen	The most common form, of 150 kDa, and the less common form of 43 kDa. Factor H is secreted by the liver into the blood serum. It is important in regulating the complement pathway, preventing unnecessary inflammation which can damage the host tissue. Complement factor H functions as a cofactor in the inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic
	The most common form, of 150 kDa, and the less common form of 43 kDa. Factor H is secreted by the liver into the blood serum. It is important in regulating the complement pathway, preventing unnecessary inflammation which can damage the host tissue. Complement factor H functions as a cofactor in the inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.
Immunogen	The most common form, of 150 kDa, and the less common form of 43 kDa. Factor H is secreted by the liver into the blood serum. It is important in regulating the complement pathway, preventing unnecessary inflammation which can damage the host tissue. Complement factor H functions as a cofactor in the inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.  Purified human factor H from serum.  mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The
Immunogen Notes	The most common form, of 150 kDa, and the less common form of 43 kDa. Factor H is secreted by the liver into the blood serum. It is important in regulating the complement pathway, preventing unnecessary inflammation which can damage the host tissue. Complement factor H functions as a cofactor in the inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.  Purified human factor H from serum.  mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The
Immunogen Notes Product Application Details	The most common form, of 150 kDa, and the less common form of 43 kDa. Factor H is secreted by the liver into the blood serum. It is important in regulating the complement pathway, preventing unnecessary inflammation which can damage the host tissue. Complement factor H functions as a cofactor in the inactivation of C3b by factor I. It makes C3b susceptible to cleavage by factor I, resulting in iC3b. Factor H also inhibits the formation of the C3bBb complex (C3 convertase) and increases the rate of dissociation of both C3 convertase and the (C3b)NBB complex (C5 convertase). This prevents these components of the classical and of alternative complement pathways from forming a positive feedback loop. Mutations in factor H are associated with hemolytic-uremic syndrome, age-related macular degeneration, membranoproliferative glomerulonephritis (MPGN) type II and chronic hypocomplementemic nephropathy.  Purified human factor H from serum.  mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  Western Blot, ELISA, Flow Cytometry, Immunohistochemistry,



# **Images**

Complement Factor H Antibody (10-10) [mFluor Violet 610 SE] [NBP2-52743MFV610] - Vial of mFluor Violet 610 conjugated antibody. mFluor Violet 610 is optimally excited at 421 nm by the Violet laser (405 nm) and has an emission maximum of 613 nm.





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP2-52743MFV610

NBP2-38695PEP Complement Factor H Recombinant Protein Antigen

210-TA-005 TNF-alpha [Unconjugated]
DY4779 Complement Factor H [Biotin]

M6000B-1 IL-6 [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-52743MFV610

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

